

Valors de referència: bibliografia 1972-2002

Recopilació preparada per X. Fuentes Arderiu^{1,2}

Les referències bibliogràfiques de la relació que segueix són una selecció de les que conté la base de dades PubMed (<<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi>>) de la Biblioteca Nacional de Medicina dels Estats Units d'Amèrica. El criteri de cerca utilitzat ha sigut que en el títol de l'article hi aparegués un dels termes següents: "reference values", "reference interval" o "reference limit", i que l'any de publicació estigués comprès entre el 1972 i el 2002. D'aquests articles s'han seleccionat tots els que pertanyent a l'àmbit de les ciències de laboratori clínic, i dels seleccionats s'han triat els escrits en alemany, anglès, espanyol, francès, italià i portuguès. A més, s'han afegit els articles sobre valors de referència publicats a les revistes *Bioquímica*, de l'Associació Mexicana de Bioquímica Clínica, i *Química Clínica*, de la Societat Espanyola de Bioquímica Clínica i Patologia Molecular.

Tal com consta en la base de dades PubMed, els títols dels articles escrits entre claudàtors són la traducció a l'anglès dels títols originals en una llengua diferent a l'anglesa.

Relació d'articles per ordre cronològic decreixent

1. Lahti A, Hyltoft Petersen P, Boyd JC. Impact of subgroup prevalences on partitioning of Gaussian-distributed reference values. *Clin Chem* 2002;48:1987-99.
2. Santos EC, Camara VM, Jesus IM, Brabo ES, Loureiro EC, Mascarenhas AF, Fayal KF, Sa Filho GC, Sagica FE, Lima MO, Higuchi H, Silveira IM. A contribution to the establishment of reference values for total mercury levels in hair and fish in amazonia. *Environ Res* 2002;90:6-11.
3. Venta R, Prieto B, Alvarez FV. Regression-based reference limits for urinary amino acids in a pediatric population. *Clin Chem Lab Med* 2002;40:383-90.
4. Hamwi A, Endler G, Rubi K, Wagner O, Endler AT. Reference values for a heterogeneous ferritin assay and traceability to the 3rd International Recombinant Standard for Ferritin (NIBSC code 94/572). *Clin Chem Lab Med* 2002;40:365-70.
5. Apostoli P, Cortesi I, Mangili A, Elia G, Drago I, Gagliardi T, Soleo L, Valente T, Sciarra GF, Aprea C, Ronchi A, Minoia C. Assessment of reference values for mercury in urine: the results of an Italian polycentric study. *Sci Total Environ* 2002;289:13-24.
6. Wahlin A, Backman L, Hultdin J, Adolfsson R, Nilsson LG. Reference values for serum levels of vitamin B12 and folic acid in a population-based sample of adults between 35 and 80 years of age. *Public Health Nutr* 2002;5:505-11.
7. Hunzelmann N, Kurschat P, Hani N, Jarisch A, Mauch C. Applicability of reference values for the determination of serum S100 protein as a marker of malignant melanoma in children. *Br J Dermatol* 2002;146:536-7. [Sense resum]
8. Kaaba SA, Al Fadhli S, Khamis A. Reference values of lymphocyte subsets in the normal healthy adult Kuwaiti Arab population. *Immunol Lett* 2002;81:199-203.
9. Nussinovitch M, Prais D, Volovitz B, Amir J, Harel D, Nussinovitch A. Reference values for lactic dehydrogenase activity and isoenzyme distribution in cerebrospinal fluid in neonates with fever but no evidence of cerebral disease. *Am J Perinatol* 2002;19:109-14.
10. Apostoli P, Baj A, Bavazzano P, Ganzi A, Neri G, Ronchi A, Soleo L, Di LL, Spinelli P, Valente T, Minoia C. Blood lead reference values: the results of an Italian polycentric study. *Sci Total Environ* 2002;287:1-11.

¹ Servei de Bioquímica Clínica, Hospital Universitari de Bellvitge, l'Hospitalet de Llobregat.

² Citació recomanada: Fuentes Arderiu X, recopilador. Valors de referència: bibliografia 1972-2002. *In vitro veritas* 2003; 4: <http://www.acclc.cat/invitroveritas/vol4/art43.html>.

11. Fischbach M, Graff V, Terzic J, Bergere V, Oudet M, Hamel G. Impact of age on reference values for serum concentration of cystatin C in children. *Pediatr Nephrol* 2002;17:104-6.
12. Tornhage CJ. Reference values for morning salivary cortisol concentrations in healthy school-aged children. *J Pediatr Endocrinol Metab* 2002;15:197-204.
13. Remer T, Neubert A, Maser-Gluth C. Anthropometry-based reference values for 24-h urinary creatinine excretion during growth and their use in endocrine and nutritional research. *Am J Clin Nutr*. 2002;75:561-9.
14. Muntau AC, Streiter M, Kappler M, Roschinger W, Schmid I, Rehnert A, Schramel P, Roscher AA. Age-related reference values for serum selenium concentrations in infants and children. *Clin Chem* 2002;48:555-60.
15. Lackmann GM. Polychlorinated biphenyls and hexachlorobenzene in full-term neonates. Reference values updated. *Biol Neonate* 2002;81:82-5.
16. Takahashi M, Stanton E, Moreno JI, Jackowski G. Immunoassay for serum glutamine synthetase in serum: development, reference values, and preliminary study in dementias. *Clin Chem* 2002;48:375-8. [Sense resum]
17. Lahti A, Hyltoft Petersen P, Boyd JC, Fraser CG, Jorgensen N. Objective criteria for partitioning gaussian-distributed reference values into subgroups. *Clin Chem* 2002;48:338-52.
18. Schiele F, Vincent-Viry M, Starck M, Beaud B, Hennache G, Siest G, Visvikis S, Herbeth B. Apolipoprotein E in apolipoprotein B (apo B)- and non-apo B-containing lipoproteins in 3523 participants in the Stanislas cohort: biological variation and genotype-specific reference limits. *Clin Chem* 2002;48:291-300.
19. Imai T, Takase M, Takeda S, Kougo T. Serum KL-6 levels in pediatric patients: reference values for children and levels in pneumonia, asthma, and measles patients. *Pediatr Pulmonol* 2002;33:135-41.
20. Armstrong GP, Barker AN, Patel H, Hart HH. Reference interval for troponin I on the ACS:Centaur assay: a recommendation based on the recent redefinition of myocardial infarction. *Clin Chem* 2002;48:198-9. [Sense resum]
21. Uhlmann EJ, Hock KG, Issitt C, Sneeringer MR, Cervelli DR, Gorman RT, Scott MG. Reference intervals for plasma cystatin C in healthy volunteers and renal patients, as measured by the Dade Behring BN II System, and correlation with creatinine. *Clin Chem* 2001;47:2031-3. [Sense resum]
22. Morison IM, Ferguson EL. Reference limits for haemoglobin and ferritin. Differences in haemoglobin concentrations reflect physiological differences. *Brit Med J* 2001;323:807-8. [Sense resum]
23. Heath AL, Fairweather-Tait S, Worwood M. Reference limits for haemoglobin and ferritin. If it's not broken, don't fix it. *Brit Med J* 2001;323:806-7 (discussion 807-8). [Sense resum]
24. [No authors listed] Clinical and professional reference guides. Table of reference intervals. *MLO Med Lab Obs* 2001;33(8 Suppl):14. [Sense resum]
25. Kost GJ. Clinical and professional reference guides. Table of critical limits. *MLO Med Lab Obs* 2001;33(8 Suppl):12-3. [Sense resum]
26. Sjogren M, Vanderstichele H, Agren H, Zachrisson O, Edsbacke M, Wikkelso C, Skoog I, Wallin A, Wahlund LO, Marcusson J, Nagga K, Andreasen N, Davidsson P, Vanmechelen E, Blennow K. Tau and Abeta42 in cerebrospinal fluid from healthy adults 21-93 years of age: establishment of reference values. *Clin Chem* 2001;47:1776-81.
27. O'Brien JE, Brookes M. Determination of reference values for a novel ketoamine-specific fructosamine assay for assessment of diabetic glycemic control. *Diabetes Technol Ther* 1999 Winter;1:447-55.

28. Henny J, Petitclerc C, Fuentes-Arderiu X, Petersen PH, Queralto JM, Schiele F, Siest G. [Revising the concept of reference values: a necessity]. *Ann Biol Clin* 2001;59:383-92. [Revisió] [Francès] [Sense resum]
29. Van den Bosch G, Van den Bossche J, Wagner C, De Schouwer P, Van De Vyvere M, Neels H. Determination of iron metabolism-related reference values in a healthy adult population. *Clin Chem* 2001;47:1465-7. [Sense resum]
30. Lippi G, Manzato F, Franchini M, Brocco G, Florenziani G, Guidi G. Establishment of reference values for the PFA-100 platelet function analyzer in pediatrics. *Clin Exp Med* 2001;1:69-70. [Sense resum]
31. Suominen P, Irjala K. Health-related decision limits rather than reference limits to guide the use of ferritin measurements. *Clin Biochem* 2001;34:345-6. [Sense resum]
32. Nguyen HN, Matern S. [Reference values in patients with daily alcohol consumption]. *Dtsch Med Wochenschr* 2001;126:767-8. [Alemany] [Sense resum]
33. Giavarina D, Mezzena G, Dorizzi RM, Soffiati G. Reference interval of D-dimer in pregnant women. *Clin Biochem* 2001;34:331-3. [Sense resum]
34. Burc L, Guibourdenche J, Luton D, Noel M, Beaudoux JL, De Lagausie P, Oury JF, Porquet D. Establishment of reference values of five amniotic fluid enzymes. Analytical performances of the Hitachi 911. Application to complicated pregnancies. *Clin Biochem* 2001;34:317-22.
35. Salazar TE, Mericq MV, Espinoza M, Iniguez G, de Carvallo P, Cassorla F. [Reference values of growth hormone binding protein (GHBP) for a normal pediatric population]. *Rev Med Chil* 2001;129:382-9. [Espanyol]
36. Chumlea WC, Guo SS, Zeller CM, Reo NV, Baumgartner RN, Garry PJ, Wang J, Pierson RN Jr, Heymsfield SB, Siervogel RM. Total body water reference values and prediction equations for adults. *Kidney Int* 2001;59:2250-8.
37. Blain H, Lerouge S, Blain A, Lacomski D, Virion JM, Humbert JC, Jeandel C. [Determination by flow cytometry of reference values of erythrocyte parameters in aged subjects]. *Presse Med* 2001;30:779-84. [Francès]
38. Rushton DH, Dover R, Sainsbury AW, Norris MJ, Gilkes JJ, Ramsay ID. Why should women have lower reference limits for haemoglobin and ferritin concentrations than men? *Brit Med J* 2001;322:1355-7. [Sense resum]
39. Christiansen M, Hogdall CK, Andersen JR, Norgaard-Pedersen B. Alpha-fetoprotein in plasma and serum of healthy adults: preanalytical, analytical and biological sources of variation and construction of age-dependent reference intervals. *Scand J Clin Lab Invest* 2001;61:205-15.
40. Jorgensen LG, Stahl M, Brandslund I, Hyltoft Petersen P, Borch-Johnsen K, de Fine Olivarius N. Plasma glucose reference interval in a low-risk population. Impact of the new WHO and ADA recommendations on the diagnosis of diabetes mellitus. *Scand J Clin Lab Invest* 2001;61:181-90.
41. Deneys V, Mazzon AM, Marques JL, Benoit H, De Bruyere M. Reference values for peripheral blood B-lymphocyte subpopulations: a basis for multiparametric immunophenotyping of abnormal lymphocytes. *J Immunol Methods* 2001;253:23-36.
42. Chiesa C, Signore F, Assumma M, Buffone E, Tramontozzi P, Osborn JF, Pacifico L. Serial measurements of C-reactive protein and interleukin-6 in the immediate postnatal period: reference intervals and analysis of maternal and perinatal confounders. *Clin Chem* 2001;47:1016-22.
43. Kinoshita S, Toyofuku M, Iida H, Wakiyama M, Kurihara M, Nakahara M, Nakata M, Nakashima K, Seo S, Hosaka N, Yano J, Misumoto T, Ishihara H, Ikeda K, Tsuchihashi M, Kawashima H, Imoto Y, Imamura K, Urabe Y, Shinohara K, Ooishi K, Abe T, Jinnouchi J, Hyoudou K, Kondo S, Kobayashi T, Ono J, Hamasaki N. Standardization of laboratory data and establishment of reference intervals in the Fukuoka Prefecture: a Japanese perspective. *Clin Chem Lab Med* 2001;39:256-62.

44. Regeczy N, Gorog G, Paloczi K. Developing an expert system for immunophenotypical diagnosis in immunodeficiency. Age-related reference values of peripheral blood lymphocyte subpopulations in Hungary. *Immunol Lett* 2001 May 1;77:47-54.
45. Ferre-Masferrer M, Fuentes-Arderiu X, Goma-Llongueras M, Aluma-Trullas A, Aramendi-Ramos M, Castano-Vidriales JL, Esteban-Salan M, Graells-Ferrer M, Jimenez-Gonzalez A, Jimenez-Lobo C, López-Esparza M, Pares-Pollan L, Regulez-Uranga M, Riesco-Prieto M, Rodriguez-Camba A, Vidal-Martínez J. Regional reference values for some quantities measured with the ADVIA Centaur analyser. A model of co-operation between the in vitro diagnostic industry and clinical laboratories. *Clin Chem Lab Med* 2001;39:166-9.
46. Durinx C, Neels H, Van der Auwera JC, Naelaerts K, Scharpe S, De Meester I. Reference values for plasma dipeptidyl-peptidase IV activity and their association with other laboratory parameters. *Clin Chem Lab Med* 2001;39:155-9.
47. Grasa Ullrich J, Rite Gracia S, Grasa Biec J, Marco Tello A, Rite Montanes S. [Reference values of interleukin-6 and tumor necrosis factor-alpha in healthy newborn infants]. *An Esp Pediatr* 2001;54:526-7. [Espanyol] [Sense resum]
48. Suominen P, Virtanen A, Lehtonen-Veromaa M, Heinonen OJ, Salmi TT, Alanen M, Mottonen T, Rajamaki A, Irjala K. Regression-based reference limits for serum transferrin receptor in children 6 months to 16 years of age. *Clin Chem* 2001;47:935-7. [Sense resum]
49. Venta R. Year-long validation study and reference values for urinary amino acids using a reversed-phase HPLC method. *Clin Chem* 2001;47:575-83.
50. Mangwendeza MP, Mandisodza A, Siziya S. Haematology reference values for healthy elderly blacks residing in Harare, Zimbabwe. *Cent Afr J Med* 2000;46:120-3.
51. Saw S, Aw TC. Age-related reference intervals for free and total prostate-specific antigen in a Singaporean population. *Pathology* 2000;32:245-9.
52. Nja F, Roksund OD, Carlsen KH. Eosinophil cationic protein (ECP) in school children living in a mountainous area of Norway: a population-based study of ECP as a tool for diagnosing asthma in children with reference values. *Allergy* 2001;56:138-44.
53. Hansen AM, Garde AH, Christensen JM, Eller N, Knudsen LE, Heinrich-Ramm R. Reference interval and subject variation in excretion of urinary metabolites of nicotine from non-smoking healthy subjects in Denmark. *Clin Chim Acta* 2001;304:125-32.
54. Steinmetz J, Spyckerelle Y, De Talance N, Fournier B, Boulange M, Leclere J, Giordanella JP. Factors of variation and reference values for TSH in 45-70 year old women. *Ann Endocrinol* 2000;61:501-507.
55. Andrew CE, Hanning I, McBain AM, Moody D, Price A. A model for a multicentre approach to the derivation of reference intervals for thyroid hormones and testosterone for laboratories using identical analysers. *Clin Chem Lab Med* 2000;38:1013-9.
56. Chenillot O, Henny J, Steinmetz J, Herbeth B, Wagner C, Siest G. High sensitivity C-reactive protein: biological variations and reference limits. *Clin Chem Lab Med* 2000;38:1003-11.
57. Gassler N, Peuschel T, Pankau R. Pediatric reference values of estradiol, testosterone, lutropin, follitropin and prolactin. *Clin Lab* 2000;46:553-60.
58. Harmoinen A, Ylinen E, Ala-Houhala M, Janas M, Kaila M, Kouri T. Reference intervals for cystatin C in pre- and full-term infants and children. *Pediatr Nephrol* 2000;15:105-8.
59. Nilsson-Ehle H, Jagenburg R, Landahl S, Svanborg A. Blood haemoglobin declines in the elderly: implications for reference intervals from age 70 to 88. *Eur J Haematol* 2000;65:297-305.
60. Rahil-Khazen R, Bolann BJ, Ulvik RJ. Trace element reference values in serum determined by inductively coupled plasma atomic emission spectrometry. *Clin Chem Lab Med* 2000;38:765-72.

61. Lun A, Schmitt M, Renz H. Phagocytosis and oxidative burst: reference values for flow cytometric assays independent of age. *Clin Chem* 2000;46:1836-9. [Sense resum]
62. Royston P, Wright EM. Goodness-of-fit statistics for age-specific reference intervals. *Stat Med* 2000;19:2943-62.
63. Kallner A, Gustavsson E, Hendig E. Can age and sex related reference intervals be derived for non-healthy and non-diseased individuals from results of measurements in primary health care? *Clin Chem Lab Med* 2000;38:633-54.
64. Henny J, Petittlerc C, Fuentes-Arderiu X, Petersen PH, Queralto JM, Schiele F, Siest G. Need for revisiting the concept of reference values. *Clin Chem Lab Med* 2000;38:589-95.
65. Orisaka M, Koyamada K, Saika Y, Sakurabayashi I. Reference values of plasma lipoprotein (a) in newly classified apolipoprotein(a) phenotype groups in the Japanese population. *Ann Clin Biochem* 2000;37:701-7.
66. Cortes-Blanco A, Ferrandez Longas A, Mayayo Dehesa E, Labarta Aizpun JI, Martínez-Lazaro R. [Hormonal reference values for adrenocortical function in healthy children from Zaragoza]. *An Esp Pediatr* 2000;52:106-15. [Espanyol]
67. Pannek J, Brands FH. [Additional aids in detection of prostate carcinomas? PSA-prostatic volume quotient, PSA-doubling time, age-dependent PSA reference values and PSA in urine]. *Urologe A* 2000;39:324-9. [Revisió] [Alemany]
68. Garcia MJ, Alegria A, Barbera R, Farre R, Lagarda MJ. Selenium, copper, and zinc indices of nutritional status: influence of sex and season on reference values. *Biol Trace Elem Res* 2000;73:77-83.
69. Theobald S. [New reference values for nutritional requirements]. *Med Monatsschr Pharm* 2000;23:232-6. [Alemany] [Sense resum]
70. Gonzalbez J, Villabona C, Ramon J, Navarro MA, Gimenez O, Ricart W, Soler J. Establishment of reference values for standard dose short synacthen test (250 microgram), low dose short synacthen test (1 microgram) and insulin tolerance test for assessment of the hypothalamo-pituitary-adrenal axis in normal subjects. *Clin Endocrinol* 2000;53:199-204.
71. Wong M, Schlaggar BL, Buller RS, Storch GA, Landt M. Cerebrospinal fluid protein concentration in pediatric patients: defining clinically relevant reference values. *Arch Pediatr Adolesc Med* 2000;154:827-31.
72. Gonzalez-Sagrado M, Martin-Gil FJ, López-Hernandez S, Fernandez-Garcia N, Olmos-Linares A, Arranz-Pena ML. Reference values and methods comparison of a new testosterone assay on the AxSYM system. *Clin Biochem* 2000;33:175-9.
73. Manahan KJ, Fanning J. Peritoneal fluid urea nitrogen and creatinine reference values. *Obstet Gynecol* 1999;93:780-2.
74. Jou HJ, Shyu MK, Shih JC, Chang MY, Lim CC, Tzeng CY, Chen SM, Hsieh FJ. Second trimester maternal serum hCG level in an Asian population: normal reference values by ultrasound dating. *J Matern Fetal Med* 2000;9:118-21.
75. Haschke-Becher E, Baumgartner M, Bachmann C. Assay of D-lactate in urine of infants and children with reference values taking into account data below detection limit. *Clin Chim Acta* 2000;298:99-109.
76. Bussieres L, Souberbielle JC, Pinto G, Adan L, Noel M, Brauner R. The use of insulin-like growth factor 1 reference values for the diagnosis of growth hormone deficiency in prepubertal children. *Clin Endocrinol* 2000;52:735-9.
77. Linnet K. Nonparametric estimation of reference intervals by simple and bootstrap-based procedures. *Clin Chem* 2000;46:867-9. [Sense resum]
78. Senofonte, Violante N, Caroli S. Assessment of reference values for elements in human hair of urban schoolboys. *J Trace Elem Med Biol* 2000;14:6-13.

79. Donnelly JG, Isotalo PA. Non-fasting reference intervals for the Abbott IMx homocysteine and AxSYM plasma folate assays: influence of the methylenetetrahydrofolate reductase 677 C-->T mutation on homocysteine. *Ann Clin Biochem* 2000;37:390-8.
80. Komaromy-Hiller G, Ash KO, Costa R, Howerton K. Comparison of representative ranges based on U.S. patient population and literature reference intervals for urinary trace elements. *Clin Chim Acta* 2000;296:71-90.
81. Chiesa ME, Fink NE. Reference intervals for erythropoietin measurement by the JCL EIA assay need to be determined locally. *Clin Chem Lab Med* 2000;38:65-6. [Sense resum]
82. Imoberdorf R, Ballmer PE. [Reference: Egger M. Under the metascope: possibilities and limits of met-analyses. *Schweiz Med Wochenschr* 1998;128:1893-901]. *Schweiz Med Wochenschr* 2000;130:410. [Alemany] [Sense resum]
83. Cuer M, Barnier A, de La Salmoniere P, Durand G, Seta N. Fluorimetric measurement of plasma alpha-L-fucosidase activity with a centrifugal analyzer: reference values in a healthy French adult population. *Clin Chem* 2000;46:562-4. [Sense resum]
84. Hjorth L, Helin I, Grubb A. Age-related reference limits for urine levels of albumin, orosomucoid, immunoglobulin G and protein HC in children. *Scand J Clin Lab Invest* 2000;60:65-73.
85. Erlandsen EJ, Randers E. Reference interval for serum C-reactive protein in healthy blood donors using the Dade Behring N Latex CRP mono assay. *Scand J Clin Lab Invest* 2000;60:37-43.
86. Indrayan A, Satyanarayana L. Reference values in medicine and validity of diagnostic test. *Indian Pediatr* 2000;37:285-91. [Revisió] [Sense resum]
87. Zardo L, Secchiero S, Sciacovelli L, Bonvicini P, Plebani M. Reference intervals: are interlaboratory differences appropriate? *Clin Chem Lab Med* 1999;37:1131-3. [Sense resum]
88. Kuriyama M, Uno H, Watanabe H, Yamanaka H, Saito Y, Shida K. Determination of reference values for total PSA, F/T and PSAD according to prostatic volume in japanese prostate cancer patients with slightly elevated serum PSA levels. *Jpn J Clin Oncol* 1999;29:617-22.
89. Conway AM, Hinchliffe RF, Finn A, Bell F. Reference values for the activated partial thromboplastin time in infants using a synthetic reagent. *Clin Lab Haematol* 1999;21:427-8. [Sense resum]
90. Dorizzi RM, Fortunato A, Marchi G, Scattolo N. Reference interval of ferritin in premenopausal women calculated in four laboratories using three different analyzers. *Clin Biochem* 2000;33:75-7. [Sense resum]
91. Cortes-Blanco A, Mayayo Dehesa E, Ferrandez Longas A, Labarta Aizpun JI, Martínez-Lazaro R. [Reference values for thyroid hormones, thyrotropin and thyroglobulin in healthy children of Zaragoza]. *An Esp Pediatr* 1999;51:361-8. [Espanyol]
92. Alimonti A, Petrucci F, Laurenti F, Papoff P, Caroli S. Reference values for selected trace elements in serum of term newborns from the urban area of Rome. *Clin Chim Acta* 2000;292:163-73.
93. Chevenne D, Trivin F, Porquet D. Insulin assays and reference values. *Diabetes Metab* 1999;25:459-76. [Revisió]
94. Wu TL, Chang CP, Tsao KC, Sun CF, Wu JT. Development of a microplate assay for serum chromogranin A (CgA): establishment of normal reference values and detection of elevated CgA in malignant diseases. *J Clin Lab Anal* 1999;13:312-9.
95. Mathioudakis B. Recommendations for vitamin intake in the European Union and the use of reference values in existing and future community legislation. *Bibl Nutr Dieta* 2001;(55):14-21. [Revisió] [Sense resum]
96. Petrini C, Alessio MG, Scapellato L, Brambilla S, Franzini C. Serum proteins by capillary zone electrophoresis: approaches to the definition of reference values. *Clin Chem Lab Med* 1999;37:975-80.
97. Fitch PS, Brown V, Schock BC, Taylor R, Ennis M, Shields MD. Serum eosinophil cationic protein (ECP): reference values in healthy nonatopic children. *Allergy* 1999;54:1199-203.

98. Cataldi L, Mussap M, Bertelli L, Ruzzante N, Fanos V, Plebani M. Cystatin C in healthy women at term pregnancy and in their infant newborns: relationship between maternal and neonatal serum levels and reference values. *Am J Perinatol* 1999;16:287-95.
99. Horn PS, Pesce AJ, Copeland BE. Reference interval computation using robust vs parametric and nonparametric analyses. *Clin Chem* 1999;45:2284-5. [Sense resum]
100. [No authors listed] Table of reference intervals. *MLO Med Lab Obs* 1999;Suppl:10. [Sense resum]
101. Lotz J, Hafner G, Prellwitz W. Reference values for a homogeneous Ferritin assay and traceability to the 3rd International Recombinant Standard for Ferritin (NIBSC code 94/572). *Clin Chem Lab Med* 1999;37:821-5.
102. Torra M, Rodamilans M, Corbella J, Ferrer R, Mazzara R. Blood chromium determination in assessing reference values in an unexposed Mediterranean population. *Biol Trace Elem Res* 1999;70:183-9.
103. Ishiguro A, Nakahata T, Matsubara K, Hayashi Y, Kato T, Suzuki Y, Shimbo T. Age-related changes in thrombopoietin in children: reference interval for serum thrombopoietin levels. *Br J Haematol* 1999;106:884-8.
104. Lun A, Ziebig R, Hammer H, Otting U, Filler G, Sinha P. Reference values for neonates and children for the UF-100 urine flow cytometer. *Clin Chem* 1999;45:1879-80. [Sense resum]
105. Randers E, Krue S, Erlandsen EJ, Danielsen H, Hansen LG. Reference interval for serum cystatin C in children. *Clin Chem* 1999;45:1856-8. [Sense resum]
106. Wright EM, Royston P. Calculating reference intervals for laboratory measurements. *Stat Methods Med Res* 1999;8:93-112. [Revisió]
107. Cortes Blanco A, Labarta Aizpun JI, Ferrandez Longas A, Mayayo Dehesa E. [Reference values for IGF-I, IGFBP-1, IGFBP-3 and osteocalcin in healthy children in Zaragoza]. *An Esp Pediatr* 1999;51:167-74. [Espanyol]
108. Cortes Blanco A, Ferrandez Longas A, Mayayo Dehesa E, Labarta Aizpun JI. [Reference values of FSH, LH, total testosterone, free testosterone, 17-beta-estradiol and SHBG in healthy children in Zaragoza]. *An Esp Pediatr* 1999;51:159-66. [Espanyol]
109. Le Moli R, Endert E, Fliers E, Mulder T, Prummel MF, Romijn JA, Wiersinga WM. Establishment of reference values for endocrine tests. II: Hyperprolactinemia. *Neth J Med* 1999;55:71-5.
110. Bona G, Petri A, Rapa A, Conti A, Sartorio A. The impact of gender, puberty and body mass on reference values for urinary growth hormone (GH) excretion in normally growing non-obese and obese children. *Clin Endocrinol* 1999;50:775-81.
111. Bokenkamp A, Dieterich C, Schumann G. Pediatric reference values for cystatin C revisited. *Pediatr Nephrol* 1999;13:367-8. [Sense resum]
112. Wudy SA, Dorr HG, Solleder C, Djalali M, Homoki J. OMIM Profiling steroid hormones in amniotic fluid of midpregnancy by routine stable isotope dilution/gas chromatography-mass spectrometry: reference values and concentrations in fetuses at risk for 21-hydroxylase deficiency. *J Clin Endocrinol Metab* 1999;84:2724-8.
113. Back SE, Nilsson JE, Fex G, Jeppson JO, Rosen U, Tryding N, von Schenck H, Norlund L. Towards common reference intervals in clinical chemistry. An attempt at harmonization between three hospital laboratories in Skane, Sweden. *Clin Chem Lab Med* 1999;37:573-92.
114. Ahokoski O, Virtanen A, Kairisto V, Scheinin H, Huupponen R, Irjala K. Biological day-to-day variation and reference change limits of serum cortisol and aldosterone in healthy young men on unrestricted diets. *Clin Chem* 1999;45:1097-9. [Sense resum]
115. Zurakowski D, Di Canzio J, Majzoub JA. Pediatric reference intervals for serum thyroxine, triiodothyronine, thyrotropin, and free thyroxine. *Clin Chem* 1999;45:1087-91. [Sense resum]

116. Dexter F, Penning DH, Lubarsky DA, DeLong E, Sanderson I, Gilbert BC, Bell E, Reves JG. Use of an automated anesthesia information system to determine reference limits for vital signs during cesarean section. *J Clin Monit Comput* 1998;14:491-8.
117. Rossi E, Beilby JP, McQuillan BM, Hung J. Biological variability and reference intervals for total plasma homocysteine. *Ann Clin Biochem* 1999;36:56-61.
118. McNerlan SE, Alexander HD, Rea IM. Age-related reference intervals for lymphocyte subsets in whole blood of healthy individuals. *Scand J Clin Lab Invest* 1999;59:89-92.
119. MacLean DR, Petrasovits A, Connelly PW, Joffres M, O'Connor B, Little JA. Plasma lipids and lipoprotein reference values, and the prevalence of dyslipoproteinemia in Canadian adults. Canadian Heart Health Surveys Research Group. *Can J Cardiol* 1999;15:434-44.
120. Connelly PW, Poapst M, Davignon J, Lussier-Cacan S, Reeder B, Lessard R, Hegele RA, Csima A. Reference values of plasma apolipoproteins A-I and B, and association with nonlipid risk factors in the populations of two Canadian provinces: Quebec and Saskatchewan. Canadian Heart Health Surveys Research Group. *Can J Cardiol* 1999;15:409-18.
121. Sibille M, Deigat N, Durieu I, Guillaumont M, Morel D, Bienvenu J, Massignon D, Durand DV. Laboratory data in healthy volunteers: reference values, reference changes, screening and laboratory adverse event limits in Phase I clinical trials. *Eur J Clin Pharmacol* 1999;55:13-9.
122. Sallam HN, Sallam A, Ezzeldin F, Agamia AF, Abou-Ali A. Reference values for the midluteal plasma progesterone concentration: evidence from human menopausal gonadotropin-stimulated pregnancy cycles. *Fertil Steril* 1999;71:711-4.
123. Ezeamuzie CI, Al-Ali SF, Al-Dowaisan A, Khan M, Hijazi Z, Thomson MS. Reference values of total serum IgE and their significance in the diagnosis of allergy among the young adult Kuwaiti population. *Clin Exp Allergy* 1999;29:375-81.
124. Flegar-Mestric Z, Nazor A, Jagarinec N. Reference intervals for haematological parameters in urban school children and adolescents. *Clin Lab Haematol* 1999;21:72-4. [Sense resum]
125. Aprea C, Betta A, Catenacci G, Lotti A, Magnaghi S, Barisano A, Passini V, Pavan I, Sciarra G, Vitalone V, Minoia C. Reference values of urinary 3,5,6-trichloro-2-pyridinol in the Italian population-validation of analytical method and preliminary results (multicentric study). *J AOAC Int* 1999;82:305-12.
126. Guillen M, Corella D, Cabello ML, Garcia AM, Hernandez-Yago J. Reference values of urinary excretion of cystine and dibasic aminoacids: classification of patients with cystinuria in the Valencian Community, Spain. *Clin Biochem* 1999;32:25-30.
127. Ferre-Masferrer M, Fuentes-Arderiu X, Puchal-Ane R. Indirect reference limits estimated from patients' results by three mathematical procedures. *Clin Chim Acta* 1999;279:97-105.
128. Baraldi E, Azzolin NM, Cracco A, Zacchello F. Reference values of exhaled nitric oxide for healthy children 6-15 years old. *Pediatr Pulmonol* 1999;27:54-8.
129. Berth M, Delanghe J, Langlois M, De Buyzere M. Reference values of serum IgA subclasses in caucasian adults by immunonephelometry. *Clin Chem* 1999;45:309-10. [Sense resum]
130. Gustafsson O, Mansour E, Norming U, Carlsson A, Tornblom M, Nyman CR. Prostate-specific antigen (PSA), PSA density and age-adjusted PSA reference values in screening for prostate cancer--a study of a randomly selected population of 2,400 men. *Scand J Urol Nephrol* 1998;32:373-7.
131. Kluge M, Riedl S, Erhart-Hofmann B, Hartmann J, Waldhauser F. Improved extraction procedure and RIA for determination of arginine8-vasopressin in plasma: role of premeasurement sample treatment and reference values in children. *Clin Chem* 1999;45:98-103.
132. Vestergaard EM, Hein HO, Meyer H, Grunnet N, Jorgensen J, Wolf H, Orntoft TF. Reference values and biological variation for tumor marker CA 19-9 in serum for different Lewis and secretor genotypes and evaluation of secretor and Lewis genotyping in a Caucasian population. *Clin Chem* 1999;45:54-61.

133. Sonntag J, Brandenburg U, Polzehl D, Strauss E, Vogel M, Dudenhausen JW, Obladen M. Complement system in healthy term newborns: reference values in umbilical cord blood. *Pediatr Dev Pathol* 1998;1:131-5.
134. Jensen NM, Brandsborg M, Boesen AM, Yde H, Dahlerup JF. Low-dose oral iron absorption test: establishment of a reference interval. *Scand J Clin Lab Invest* 1998;58:511-9.
135. Kroll J, Saxtrup O. On the use of patient data for the definition of reference intervals in clinical chemistry. *Scand J Clin Lab Invest* 1998;58:469-73.
136. de Bos Kuil MJ, Endert E, Fliers E, Prummel MF, Romijn JA, Wiersinga WM. Establishment of reference values for endocrine tests. I: Cushing's syndrome. *Neth J Med* 1998;53:153-63.
137. Virtanen A, Kairisto V, Uusipaikka E. Regression-based reference limits: determination of sufficient sample size. *Clin Chem* 1998;44:2353-8.
138. Diaz J, Serrano E, Acosta F, Carbonell LF. Reference intervals for four biochemistry analytes in plasma for evaluating oxidative stress and lipid peroxidation in human plasma. *Clin Chem* 1998;44:2215-7. [Sense resum]
139. Diaz J, Fernandez-Arjona M, Perandones C, Serrano E, Frau C, Cortes I, Luque A, Carbonell LF. Reference intervals for biochemistry parameters for evaluation of oxidative stress in human sperm. *Clin Chem* 1998;44:2197-9. [Sense resum]
140. Foulon T, Laporte F, Payen N, Gros Lambert P. [Plasma apolipoproteins AI and B: reference values in the Isere population and determination of individual values]. *Ann Biol Clin* 1998;56:451-6. [Francès]
141. Diaz J, Serrano E, Acosta F, Carbonell LF. Reference interval for human plasma nitric oxide end products. *Clin Biochem* 1998;31:513-5. [Sense resum]
142. Lynch M, Thompson A, Gaumont B, Kilroy J, Leonard H, Jacob A, Jackson T. Transference of reference intervals in the validation of automated chemiluminescent immunoassays on a new platform. *Clin Chem* 1998;44:2061-3. [Sense resum]
143. Erlandsen EJ, Randers E, Kristensen JH. Reference intervals for serum cystatin C and serum creatinine in adults. *Clin Chem Lab Med* 1998;36:393-7.
144. Balsells D, Gella FJ, Gubern G, Canalias F. Reference values for alpha-amylase in human serum and urine using 2-chloro-4-nitrophenyl-alpha-D-maltotriose as substrate. *Clin Chim Acta* 1998;274:213-7. [Sense resum]
145. Lackmann GM, Schnieder C, Bohner J. Gestational age-dependent reference values for iron and selected proteins of iron metabolism in serum of premature human neonates. *Biol Neonate* 1998;74:208-13.
146. Parentoni LS, de Faria EC, Bartelega MJ, Moda VM, Facin AC, Castilho LN. Glycated hemoglobin reference limits obtained by high performance liquid chromatography in adults and pregnant women. *Clin Chim Acta* 1998;274:105-9. [Sense resum]
147. Jagarinec N, Flegar-Mestric Z, Surina B, Vrhovski-Hebrang D, Preden-Kerekovic V. Pediatric reference intervals for 34 biochemical analytes in urban school children and adolescents. *Clin Chem Lab Med* 1998;36:327-37.
148. Roberts WL, Paulson WD. Method-specific reference intervals for serum anion gap and osmolality. *Clin Chem* 1998;44:1582. [Sense resum]
149. Rivero-Marcotegui A, Olivera-Olmedo JE, Valverde-Visus FS, Palacios-Sarrasqueta M, Grijalba-Uche A, Garcia-Merlo S. Water, fat, nitrogen, and sugar content in feces: reference intervals in children. *Clin Chem* 1998;44:1540-4.
150. Simopoulos AP. Redefining dietary reference values and food safety. *World Rev Nutr Diet* 1998;83:219-22. [Revisió] [Sense resum]
151. Diaz J, Serrano E, Acosta F, Carbonell LF. Lipoperoxides kit evaluated for measuring lipoperoxides in biological samples: reference intervals for human plasma. *Clin Biochem* 1998;31:277-9. [Sense resum]

152. Takubo T, Tatsumi N, Tsuda I, Kasuya K, Taniguchi Y, Fujita T, Uchida K, Matsuo Y, Hayashi T. Reference values for bisphosphoglycerate mutase protein content and specific activity in human erythrocytes. *Acta Haematol* 1998;99:234-5. [Sense resum]
153. Boulat O, Krieg MA, Janin B, Burckhardt P, Francioli P, Bachmann C. Clinical chemistry variables in normal elderly and healthy ambulatory populations: comparison with reference values. *Clin Chim Acta* 1998;272:127-35.
154. Kintzel K, Sonntag J, Strauss E, Obladen M. Neuron-specific enolase: reference values in cord blood. *Clin Chem Lab Med* 1998;36:245-7.
155. de Siqueira ME, Martins I, Costa AC, Andrade EL, Esteves MT, Lima SA. [Reference values for carboxyhemoglobin]. *Rev Saude Publica* 1997;31:618-23. [Portuguès]
156. Blohm ME, Vesterling-Horner D, Calaminus G, Gobel U. Alpha 1-fetoprotein (AFP) reference values in infants up to 2 years of age. *Pediatr Hematol Oncol* 1998;15:135-42.
157. Vincent-Viry M, Schiele F, Gueguen R, Bohnet K, Visvikis S, Siest G. Biological variations and genetic reference values for apolipoprotein E serum concentrations: results from the STANISLAS cohort study. *Clin Chem* 1998;44:957-65.
158. Ooi DS, Innanen VT, Wang D, Chong GL, Donnelly JG, Arseneault JJ, Pronovost C, Wells G. Establishing reference intervals for DPC's free testosterone radioimmunoassay. *Clin Biochem* 1998;31:15-21.
159. Tsang CW, Lazarus R, Smith W, Mitchell P, Koutts J, Burnett L. Hematological indices in an older population sample: derivation of healthy reference values. *Clin Chem* 1998;44:96-101.
160. Puelles A, Jimenez M, Mateo G, Rodriguez MM, Eugui J. Conversion of apolipoproteins A-1 and B reference limits obtained by a turbidimetric procedure to other methods. *Ann Clin Biochem* 1998;35:311-6. [Sense resum]
161. Bokenkamp A, Domanetzki M, Zinck R, Schumann G, Brodehl J. Reference values for cystatin C serum concentrations in children. *Pediatr Nephrol* 1998;12:125-9.
162. Mold JW, Aspy CB, Blick KE, Lawler FH. The determination and interpretation of reference intervals for multichannel serum chemistry tests. *J Fam Pract* 1998;46:233-41.
163. Gozzo ML, Noia G, Barbaresi G, Colacicco L, Serraino MA, De Santis M, Lippa S, Calla C, Caruso A, Mancuso S, Giardina B. Reference intervals for 18 clinical chemistry analytes in fetal plasma samples between 18 and 40 weeks of pregnancy. *Clin Chem* 1998;44:683-5. [Sense resum]
164. Horn PS, Pesce AJ, Copeland BE. A robust approach to reference interval estimation and evaluation. *Clin Chem* 1998;44:622-31.
165. Paoliello MM, Gutierrez PR, Turini CA, Matsuo T, Mezzaroba L, Barbosa DS, Alvarenga AL, Carvalho SR, Figueiroa GA, Leite VG, Gutierrez AC, Nogueira KB, Inamine WA, Zavatti AM. [Reference values for lead levels in blood for the urban population]. *Rev Saude Publica* 1997;31:144-8. [Portuguès]
166. Wright EM, Royston P. Simplified estimation of age-specific reference intervals for skewed data. *Stat Med* 1997;16:2785-803.
167. Virtanen A, Kairisto V, Irjala K, Rajamaki A, Uusipaikka E. Regression-based reference limits and their reliability: example on hemoglobin during the first year of life. *Clin Chem* 1998;44:327-35.
168. Karr M, Mira M, Causer J, Earl J, Alperstein G, Wood F, Fett MJ, Coakley J. Age-specific reference intervals for plasma vitamins A, E and beta-carotene and for serum zinc, retinol-binding protein and prealbumin for Sydney children aged 9-62 months. *Int J Vitam Nutr Res* 1997;67:432-6.
169. Taimela E, Kairisto V, Koskinen P, Leino A, Irjala K. Reference intervals for serum thyrotropin, free thyroxine and free triiodothyronine in healthy adults in Finland, measured by an immunoautomate based on time-resolved fluorescence (AutoDELFI). *Eur J Clin Chem Clin Biochem* 1997;35:889-90.

170. Meissner D. [Reference values for blood and serum selenium in the Dresden area]. *Med Klin* 1997;92(Suppl 3):41-2. [Alemany]
171. Mahalingam TR, Vijayalakshmi S, Prabhu RK, Thiruvengadasami A, Murthy KS, Sen D, Mathews CK, Shanmugasundaram KR. Studies on some trace and minor elements in blood. A survey of the Kalpakkam (India) population. Part II: Reference values for plasma and red cell, and correlation with coronary risk index. *Biol Trace Elem Res* 1997;57:207-21.
172. Fuentes-Arderiu X, Ferre-Masferrer M, Alvarez-Funes V. Harris & Boyd's test for partitioning the reference values. *Eur J Clin Chem Clin Biochem* 1997;35:733. [Sense resum]
173. Ferre-Masferrer M, Fuentes-Arderiu X, Alvarez-Funes V, Guell-Miro R, Castineiras-Lacambra MJ. Multicentric reference values: shared reference limits. *Eur J Clin Chem Clin Biochem* 1997;35:715-8.
174. Armstrong NC, Paganga G, Brunner E, Miller NJ, Nanchahal K, Shipley M, Rice-Evans CA, Marmot MG, Diplock AT. Reference values for alpha-tocopherol and beta-carotene in the Whitehall II Study. *Free Radic Res* 1997;27:207-19.
175. Jensen KT, Carstens J, Ivarsen P, Pedersen EB. A new, fast and reliable radioimmunoassay of brain natriuretic peptide in human plasma. Reference values in healthy subjects and in patients with different diseases. *Scand J Clin Lab Invest* 1997;57:529-40.
176. Norlund L, Fex G, Lanke J, Von Schenck H, Nilsson JE, Leksell H, Grubb A. Reference intervals for the glomerular filtration rate and cell-proliferation markers: serum cystatin C and serum beta 2-microglobulin/cystatin C-ratio. *Scand J Clin Lab Invest* 1997;57:463-70.
177. Steinmetz J, Caces E, Couderc R, Beucler I, Legrand A, Henny J. [Reference values of apolipoproteins A1 and B. Contribution of international standardization. Travail collaboratif entre la SFBC, l'Arcol et le SFRL]. *Ann Biol Clin* 1997;55:451-4. [Francès]
178. Vulkov I. Percentile reference values of lipid risk factors of ischaemic heart disease: cholesterol, LDL-cholesterol, HDL-cholesterol and triglycerides in individuals aged 20 to 60 years in Plovdiv region. *Folia Med (Plovdiv)* 1997;39:37-45.
179. Khan FA, Dilawar M, Khan DA. Reference values of common blood chemistry analytes in healthy population of Rawalpindi-Islamabad area. *J Pak Med Assoc* 1997;47:156-9.
180. Marchese N, Canini S, Fabi L, Famularo L. Paediatric reference values for urinary catecholamine metabolites evaluated by high performance liquid chromatography and electrochemical detection. *Eur J Clin Chem Clin Biochem* 1997;35:533-7.
181. Nielsen F, Mikkelsen BB, Nielsen JB, Andersen HR, Grandjean P. Plasma malondialdehyde as biomarker for oxidative stress: reference interval and effects of life-style factors. *Clin Chem* 1997;43:1209-14.
182. Banales JL, Vazquez L, Mendoza F, Baltazares M, Raymond Y, Nava A, Selman M. On the correct determination of reference values for serum antibodies against pigeon serum antigen using a group of healthy blood donors. *Arch Med Res* 1997 Summer;28:289-91.
183. Rukgauer M, Klein J, Kruse-Jarres JD. Reference values for the trace elements copper, manganese, selenium, and zinc in the serum/plasma of children, adolescents, and adults. *J Trace Elem Med Biol* 1997;11:92-8.
184. Roberts WL, Johnson RD. The serum anion gap. Has the reference interval really fallen? *Arch Pathol Lab Med* 1997;121:568-72.
185. Olmedilla B, Granado F, Gil-Martínez E, Blanco I, Rojas-Hidalgo E. Reference values for retinol, tocopherol, and main carotenoids in serum of control and insulin-dependent diabetic Spanish subjects. *Clin Chem* 1997;43:1066-71.
186. Lofman O, Lindahl T, Varenhorst E. Age-specific reference values for serum prostate-specific antigen in a community-based population of healthy Swedish men. *Scand J Clin Lab Invest* 1997;57:225-32.

187. Comans-Bitter WM, de Groot R, van den Beemd R, Neijens HJ, Hop WC, Groeneveld K, Hooijkaas H, van Dongen JJ. Immunophenotyping of blood lymphocytes in childhood. Reference values for lymphocyte subpopulations. *J Pediatr* 1997;130:388-93.
188. Ries M, Klinge J, Rauch R. Age-related reference values for activation markers of the coagulation and fibrinolytic systems in children. *Thromb Res* 1997;85:341-4.
189. Nystrom FH, Ohman PK, Ekman BA, Osterlund MK, Karlberg BE, Arnqvist HJ. Population-based reference values for IGF-I and IGF-binding protein-1: relations with metabolic and anthropometric variables. *Eur J Endocrinol* 1997;136:165-72.
190. Ngou-Milama E, Djeki J, Akpona S, Dufillot D, Lebreton Devonne T, Mouray H. [Selection criteria for the establishment of reference values in tropical zones. Application to specific proteins: C-reactive protein, haptoglobin, transferrin in gabonese children]. *Bull Soc Pathol Exot* 1997;90:33-6. [Francès]
191. Ichihara K, Kawai T. Determination of reference intervals for 13 plasma proteins based on IFCC international reference preparation (CRM470) and NCCLS proposed guideline (C28-P, 1992): a strategy for partitioning reference individuals with validation based on multivariate analysis. *J Clin Lab Anal* 1997;11:117-24.
192. Itoh Y, Ichihara K, Kanno T, Sugawara T, Ohkubo A, Hirabayashi Y, Igarashi S, Kawano K, Iwata S, Saito K, Kawai T. Serum protein standardization project in Japan: evaluation of an IFCC reference material (RPPHS/CRM470) and establishment of reference intervals. *J Clin Lab Anal* 1997;11:39-44.
193. Winklhofer-Roob BM, van't Hof MA, Shmerling DH. Reference values for plasma concentrations of vitamin E and A and carotenoids in a Swiss population from infancy to adulthood, adjusted for seasonal influences. *Clin Chem* 1997;43:146-53.
194. Macy EM, Hayes TE, Tracy RP. Variability in the measurement of C-reactive protein in healthy subjects: implications for reference intervals and epidemiological applications. *Clin Chem* 1997;43:52-8.
195. Cevat Inal T, Tuli A, Yuregir GT. Evaluation of reference values for erythrocyte glutathione. *Clin Chim Acta* 1996;256:189-96.
196. Kraaijenhagen RJ. Reticulocyte reference values by the Bhattacharya method: results of a pilot study. *Clin Lab Haematol* 1996;18(Suppl 1):15-6. [Sense resum]
197. Tarallo P, Humbert JC, Fournier B, Mahassen P, Henny J. Reticulocytes: reference limits. *Clin Lab Haematol* 1996;18(Suppl 1):13-4.
198. Tencer J, Thysell H, Grubb A. Analysis of proteinuria: reference limits for urine excretion of albumin, protein HC, immunoglobulin G, kappa- and lambda-immunoreactivity, orosomuroid and alpha 1-antitrypsin. *Scand J Clin Lab Invest* 1996;56:691-700.
199. Alegria A, Barbera R, Clemente G, Farre R, Garcia MJ, Lagarda MJ. Selenium and glutathione peroxidase reference values in whole blood and plasma of a reference population living in Valencia, Spain. *J Trace Elem Med Biol* 1996;10:223-8.
200. Gallois Y, Vol S, Caces E, Balkau B. Distribution of fasting serum insulin measured by enzyme immunoassay in an unselected population of 4,032 individuals. Reference values according to age and sex. D.E.S.I.R. Study Group. *Donnees Epidemiologiques sur le Syndrome d'Insulino-Resistance. Diabetes Metab* 1996;22:427-31.
201. Lackmann GM. Reference values for selected enzyme activities in serum from healthy human neonates. *Clin Biochem* 1996;29:599-602. [Sense resum]
202. Greenway DC, Hindmarsh JT, Wang J, Khodadeen JA, Hebert PC. Reference interval for whole blood ionized magnesium in a healthy population and the stability of ionized magnesium under varied laboratory conditions. *Clin Biochem* 1996;29:515-20.
203. Najmey SS, Erickson EN Jr, Keil LB, DeBari VA. Antibodies to beta 2 glycoprotein I: standardized immunoassay and a reference interval for healthy controls. *Proc Assoc Am Physicians* 1996;108:467-72.

204. Uttenreuther-Fischer MM, Ziemer S, Gaedicke G. Resistance to activated protein C (APCR): reference values of APC-ratios for children. *Thromb Haemost* 1996;76:813-4. [Sense resum]
205. Banning A, Musumeci F, Penny W, Tovey JA. Reference intervals for cardiac troponin T, creatine kinase and creatine kinase-MB isoenzyme following coronary bypass graft surgery. *Ann Clin Biochem* 1996;33:561-2. [Sense resum]
206. Davis ML, Austin C, Messmer BL, Nichols WK, Bonin AP, Bennett MJ. IFCC-standardized pediatric reference intervals for 10 serum proteins using the Beckman Array 360 system. *Clin Biochem* 1996;29:489-92. [Sense resum]
207. Howard RR, Fasano CS, Frey L, Miller CH. Reference intervals of CD3, CD4, CD8, CD4/CD8, and absolute CD4 values in Asian and Non-Asian populations. *Cytometry* 1996;26:231-2.
208. Iversen OH, Roger M, Solberg HE, Wetteland P. Rising erythrocyte sedimentation rate during several years before diagnosis can be a predictive factor in 70% of renal cell carcinoma patients. The benefit of knowing subject-based reference values. *J Intern Med* 1996;240:133-41.
209. Orum O, Hansen M, Jensen CH, Sorensen HA, Jensen LB, Horslev-Petersen K, Teisner B. Procollagen type I N-terminal propeptide (PINP) as an indicator of type I collagen metabolism: ELISA development, reference interval, and hypovitaminosis D induced hyperparathyroidism. *Bone* 1996;19:157-63.
210. Hammerman-Rozenberg R, Cohen A, Ginsberg G, Maaravi Y, Ebstein RP, Stessman J. Laboratory reference values for the 70 year olds. *Isr J Med Sci* 1996;32:611-20.
211. Rabier D, Chadefaux-Vekemans B, Oury JF, Aupetit J, Bardet J, Gasquet M, Merhand E, Parvy P, Kamoun P. Gestational age-related reference values for amniotic fluid amino acids: a useful tool for prenatal diagnosis of aminoacidopathies. *Prenat Diagn* 1996;16:623-8.
212. Nuttall KL, Pingree SS, Ashwood ER. Reference intervals for 24-hour and random urine porphyrins. *Ann Clin Lab Sci* 1996;26:313-22.
213. Petersen PH, Fraser CG, Ricos C, Stockl D, Libeer JC, Baadenhuijsen H, Thienpont L. On establishment of common reference intervals in laboratory medicine. *Eur J Clin Chem Clin Biochem* 1996;34:515-6. [Sense resum]
214. Alfthan G, Neve J. Reference values for serum selenium in various areas-evaluated according to the TRACY protocol. *J Trace Elem Med Biol* 1996;10:77-87. [Revisió]
215. Jorgensen LG, Lober J, Carlsen NL, Momsen G, Hirsch FR. Serum neuron specific enolase (S-NSE) reference interval evaluation by time-resolved immunofluorometry compared with a radioimmunoassay. *Clin Chim Acta* 1996;249:77-91.
216. Minder EI, Schneider-Yin X. Age-dependent reference values of urinary porphyrins in children. *Eur J Clin Chem Clin Biochem* 1996;34:439-43.
217. Gilsanz F, Gomez-Tabera C, Torres M, Barredo M, Estenoz J, López-Carbajo A, Millan I. [Reference values in the diagnosis of anemia in the elderly]. *Rev Clin Esp* 1996;196:289-92. [Espanyol]
218. Rizzo V, Anesi A, Montalbetti L, Bellantoni G, Trotti R, Melzi d'Eril GV. Reference values of neuroactive amino acids in the cerebrospinal fluid by high-performance liquid chromatography with electrochemical and fluorescence detection. *J Chromatogr A* 1996;729:181-8.
219. Nagao AT, Costa-Carvalho BT, Sol D, Naspitz C, Pereira AB. Salivary secretory IgA reference values in Brazilian healthy children. *J Trop Pediatr* 1996;42:119. [Sense resum]
220. Akdag R, Energin VM, Kalayci AG, Karakelleoglu C. Reference limits for routine haematological measurements in 7-14-year-old children living at an intermediate altitude (1869 m, Erzurum, Turkey). *Scand J Clin Lab Invest* 1996;56:103-9.
221. Rasmussen K, Moller J, Lyngbak M, Pedersen AM, Dybkjaer L. Age- and gender-specific reference intervals for total homocysteine and methylmalonic acid in plasma before and after vitamin supplementation. *Clin Chem* 1996;42:630-6.

222. Contois JH, McNamara JR, Lammi-Keefe CJ, Wilson PW, Massov T, Schaefer EJ. Reference intervals for plasma apolipoprotein B determined with a standardized commercial immunoturbidimetric assay: results from the Framingham Offspring Study. *Clin Chem* 1996;42:515-23.
223. Contois J, McNamara JR, Lammi-Keefe C, Wilson PW, Massov T, Schaefer EJ. Reference intervals for plasma apolipoprotein A-1 determined with a standardized commercial immunoturbidimetric assay: results from the Framingham Offspring Study. *Clin Chem* 1996;42:507-14.
224. Borque de Larrea L. [New standardization of the measurement of serum proteins. New reference intervals]. *Med Clin* 1996;106:238-9. [Espanyol] [Sense resum]
225. Fuentes Arderiu X, Ferre-Masferrer M. On the reliability of published reference limits. *Eur J Clin Chem Clin Biochem* 1996;34:143. [Sense resum]
226. Ruxton CH, Kirk TR, Belton NR, Holmes MA. Energy and nutrient intakes in a sample of 136 Edinburgh 7-8 year olds: a comparison with United Kingdom dietary reference values. *Br J Nutr* 1996;75:151-60.
227. Ishiguro A, Inoue K, Nakahata T, Nishihira H, Kojima S, Ueda K, Suzuki Y, Shimbo T. Reference intervals for serum granulocyte colony-stimulating factor levels in children. *J Pediatr* 1996;128:208-12.
228. Cals MJ, Bories PN, Blonde-Cynober F, Coudray-Lucas C, Desveaux N, Devanlay M, Duche JC, Luciani L, Meneguzzer E, Ponteziere C, Succari M, Tortrat D. [Reference intervals and biological profile in a group of healthy elderly population in the Paris region]. *Ann Biol Clin* 1996;54:307-15. [Francès]
229. Chapuis-Cellier C, Cloppet H, Chazaud A, Coquelin H. [Immunonephelometric analysis of immunoglobulin light chains: agreement between the theoretical and experimental model. Assessment of the reference values of the kappa/lambda ratio and the heavy to light chain ratio]. *J Pharm Belg* 1996;51:23-27. [Francès]
230. Ichihara K, Kawai T. Determination of reference intervals for 13 plasma proteins based on IFCC international reference preparation (CRM470) and NCCLS proposed guideline (C28-P,1992): trial to select reference individuals by results of screening tests and application of maximal likelihood method. *J Clin Lab Anal* 1996;10:110-7.
231. Blennow K, Skoog I, Wallin A, Wikkelso C, Fredman P. Immunoglobulin M in cerebrospinal fluid: reference values derived from 111 healthy individuals 18-88 years of age. *Eur Neurol* 1996;36:201-5.
232. Bon GG, Kenemans P, Verstraeten R, van Kamp GJ, Hilgers J. Serum tumor marker immunoassays in gynecologic oncology: establishment of reference values. *Am J Obstet Gynecol* 1996;174:107-14.
233. Stulnig T, Maczek C, Bock G, Majdic O, Wick G. Reference intervals for human peripheral blood lymphocyte subpopulations from 'healthy' young and aged subjects. *Int Arch Allergy Immunol* 1995;108:205-10.
234. Sone T, Miyake M, Takeda N, Fukunaga M. Urinary excretion of type I collagen crosslinked N-telopeptides in healthy Japanese adults: age- and sex-related changes and reference limits. *Bone* 1995;17:335-9.
235. Marshall MD, Kales SN, Christiani DC, Goldman RH. Are reference intervals for carboxyhemoglobin appropriate? A survey of Boston area laboratories. *Clin Chem* 1995;41:1434-8.
236. Ducharme GR, Gannoun A, Guertin MC, Jequier JC. Reference values obtained by kernel-based estimation of quantile regressions. *Biometrics* 1995;51:1105-16.
237. Cembrowski GS, Fairbanks VF. Can hematology reference intervals be derived from hospitalized patients' data? *Clin Chem* 1995;41:1048-51. [Sense resum]
238. Diaz J, Tornel PL, Martínez P. Reference intervals for blood ammonia in healthy subjects, determined by microdiffusion. *Clin Chem* 1995;41:1048. [Sense resum]

239. López García A, Galindo García A, Paz Martínez D, Aguirre Langle E, Perez Fernandez S, Teran Vazquez L. [Reference values of T-lymphocyte subsets in umbilical cord blood]. *Rev Alerg Mex* 1995;42:60-3. [Espanyol]
240. Steinmetz J, Tarallo P, Fournier B, Caces E, Siest G. Reference limits of apolipoprotein A-I and apolipoprotein B using an IFCC standardized immunonephelometric method. *Eur J Clin Chem Clin Biochem* 1995;33:337-42.
241. Leino A, Impivaara O, Irjala K, Maki J, Peltola O, Jarvisalo J. Health-based reference intervals for ALAT, ASAT and GT in serum, measured according to the recommendations of the European Committee for Clinical Laboratory Standards (ECCLS). *Scand J Clin Lab Invest* 1995;55:243-50.
242. Negri M, Gheller A, Zanini M, Tonni S, Caputo M. Reference interval for serum secretory IgA. *Clin Chem* 1995;41:634. [Sense resum]
243. Romppanen J, Mononen I. Age-related reference values for urinary excretion of sialic acid and deoxysialic acid: application to diagnosis of storage disorders of free sialic acid. *Clin Chem* 1995;41:544-7.
244. Siest G, Galteau MM, Henny J. [Reference values of laboratory tests: a useful and important epidemiologic contribution of the Centers for health tests]. *Bull Acad Natl Med* 1995;179:235-45; discussion 245-8. [Francès]
245. Apostoli P. [Reference values for trace elements in biological fluids]. *Ann Ist Super Sanita* 1995;31:261-74. [Revisió] [Italià]
246. Zolezzi C, Lenzi L, Salizzoni MP, Pizzoferrato A. [Reference values of assayable pyridinolines as markers of bone turn-over]. *Minerva Med* 1995;86:5-9. [Italià]
247. Tanasijevic MJ, Ginsburg E, Wybenga DR, Barbieri RL, Winkelman JW. Total serum calcium reference intervals in postmenopausal outpatients. *Gynecol Obstet Invest* 1995;39:266-70.
248. Solberg HE. Subject-based reference values. *Scand J Clin Lab Invest* 1995;222(Suppl):7-10.
249. Kouri T, Kairisto V, Virtanen A, Uusipaikka E, Koivula T, Nanto V. Reliability and adequacy of discharge diagnosis databases in the production of reference values. *Scand J Clin Lab Invest* 1995;222(Suppl.):69-73.
250. Kairisto V, Kouri T, Virtanen A, Uusipaikka E, Koivula T, Nanto. Estimation of reference change limits using patient data. *Scand J Clin Lab Invest* 1995;222(Suppl):37-41.
251. Koivula T. Possibilities for quality assurance of reference intervals. *Scand J Clin Lab Invest* 1995;222(Suppl):17-20.
252. Stepniewski M, Cieszkowska U, Pach J, Kitlinski M. Reference values for fetal hemoglobin concentration in blood of steel-mill workers. *Przegl Lek* 1995;52:257-9.
253. Weykamp CW, Martina WV, van der Dijs FP, Penders TJ, van der Slik W, Muskiet FA. Hemoglobins S and C: reference values for glycohemoglobin in heterozygous, double-heterozygous and homozygous subjects, as established by 13 methods. *Clin Chim Acta* 1994;231:161-71.
254. Sokoll LJ, Russell RM, Sadowski JA, Morrow FD. Establishment of creatinine clearance reference values for older women. *Clin Chem* 1994;40:2276-81.
255. Kouri T, Kairisto V, Virtanen A, Uusipaikka E, Rajamaki A, Finneman H, Juva K, Koivula T, Nanto V. Reference intervals developed from data for hospitalized patients: computerized method based on combination of laboratory and diagnostic data. *Clin Chem* 1994;40:2209-15.
256. Solberg HE. Using a hospitalized population to establish reference intervals: pros and cons. *Clin Chem* 1994;40:2205-6. [Sense resum]
257. Boerma GJ, de Bruijn AM, van Teunenbroek A. Reference values for apolipoprotein A-I and apolipoprotein B in serum still depend on choice of assay techniques. *Eur J Clin Chem Clin Biochem* 1994;32:923-7.

258. Steinmetz J, Tarallo P, Fournier B, Jaid M, Caces E, Vol S, Tichet J, Siest G. [Reference values of lipoprotein(a) in a French population]. *Presse Med* 1994;23:1695-8. [Francès]
259. Kairisto V, Hanninen KP, Leino A, Pulkki K, Peltola O, Nanto V, Voipio-Pulkki LM, Irjala K. Generation of reference values for cardiac enzymes from hospital admission laboratory data. *Eur J Clin Chem Clin Biochem* 1994;32:789-96.
260. Mischke R, Menzel D. [Measuring fibrinogen concentrations in healthy dogs: standardization, comparison of methods and reference values]. *Zentralbl Veterinarmed A* 1994;41:587-98. [Alemany]
261. Main KM, Jarden M, Angelo L, Dinesen B, Hertel NT, Juul A, Muller J, Skakkebaek NE. The impact of gender and puberty on reference values for urinary growth hormone excretion: a study of 3 morning urine samples in 517 healthy children and adults. *J Clin Endocrinol Metab* 1994;79:865-71.
262. van der Meulen EA, Boogaard PJ, van Sittert NJ. Use of small-sample-based reference limits on a group basis. *Clin Chem* 1994;40:1698-702.
263. Liappis N, Starke A, Schlebusch H. [Comparison of the luminescence enhanced enzyme immunoassay Amerlite FT4/MAB-FT4 and Amerlite FT3/MAB-FT3 in blood serum of children. Reference values]. *Klin Padiatr* 1994;206:392-6. [Alemany]
264. Bock A, Herkner KR. Reticulocyte maturity pattern analysis as a predictive marker of erythropoiesis in paediatrics. Part I: Evaluation of age-dependent reference values. *Clin Lab Haematol* 1994;16:247-51.
265. Coates JE, Chapelsky LJ, Yatscoff RW. Pediatric reference intervals for plasma renin. *Clin Biochem* 1994;27:316-7. [Sense resum]
266. Eugui J, Logrono MJ, Ruiz R, Zugaza C, Mirabel JL, Martínez C. Immunoturbidimetry of serum apolipoproteins A-I and B on the Cobas Bio centrifugal analyzer: method validation and reference values. *Clin Biochem* 1994;27:310-5.
267. Berger J. Application of fuzzy reasoning to haematological reference values. *Nouv Rev Fr Hematol* 1994;36:333-4.
268. Kim H, Cho SH. Estimation of the geometric means and the reference values of blood lead levels among Koreans. *J Korean Med Sci* 1994;9:304-12.
269. Brown AK, Sleeper LA, Miller ST, Pegelow CH, Gill FM, Waclawiw MA. Reference values and hematologic changes from birth to 5 years in patients with sickle cell disease. Cooperative Study of Sickle Cell Disease. *Arch Pediatr Adolesc Med* 1994;148:796-804.
270. Tarallo P, Humbert JC, Mahassen P, Fournier B, Henny J. Reticulocytes: biological variations and reference limits. *Eur J Haematol* 1994;53:11-5.
271. Guneral F, Bachmann C. Age-related reference values for urinary organic acids in a healthy Turkish pediatric population. *Clin Chem* 1994;40:862-6.
272. Faulkner WR, Demers LM. Importance of age-dependent reference values in biochemical testing: are we including the elderly population? *Clin Chem* 1994;40:855-6. [Sense resum]
273. Sansores-Garcia L, Majluf-Cruz A, Vargas-Vorackova F, Labardini-Mendez J. [Reference values for protein C activity in healthy adults in Mexico]. *Rev Invest Clin* 1994;46:231-5. [Espanyol]
274. Frolich A, Nielsen BF, Nielsen J, Conradsen K, McNair P. Do local hospitals meet the analytical goals for the use of common reference intervals? *Scand J Clin Lab Invest* 1994;54:169-76.
275. Nilsson JE, Lanke J, Nilsson-Ehle P, Tryding N, Schersten B. Reference intervals and decision limits for plasma lipids and lipoproteins: a practical evaluation of current recommendations. *Scand J Clin Lab Invest* 1994;54:137-46.
276. Kogner P, Bjork O, Theodorsson E. Plasma neuropeptide Y in healthy children: influence of age, anaesthesia and the establishment of an age-adjusted reference interval. *Acta Paediatr* 1994;83:423-7.
277. Lipson SF, Ellison PT. Reference values for luteal progesterone measured by salivary radioimmunoassay. *Fertil Steril* 1994;61:448-54.

278. Farriol M, Schwartz S. Plasma carnitine reference values. *Ann Clin Biochem* 1994;31:188-9. [Sense resum]
279. Ras RM, Aulesa C, Ortega JJ. [Reference values for vitamin B12 in blood in a population of children and one of adults using the IMxRB12 method]. *Sangre* 1994;39:29-34. [Espanyol]
280. Marniemi J, Maatela J, Jarvisalo J, Reunanen A, Maki J, Tikkanen MJ. Health-based reference values of the Mini-Finland Health Survey: 3. Triglycerides in total serum and in different lipoprotein fractions. *Scand J Clin Lab Invest* 1994;54:43-50.
281. Maatela J, Marniemi J, Reunanen A, Jarvisalo J, Maki J, Tikkanen MJ. Health-based reference values of the Mini-Finland Health Survey: 2. Cholesterol in total serum and in different lipoprotein fractions. *Scand J Clin Lab Invest* 1994;54:33-42.
282. Gullestad L, Midtvedt K, Dolva LO, Norseth J, Kjekshus J. The magnesium loading test: reference values in healthy subjects. *Scand J Clin Lab Invest* 1994;54:23-31.
283. al-Tamer YY, Hadi EA. Age dependent reference intervals of glucose, urea, protein, lactate and electrolytes in thermally induced sweat. *Eur J Clin Chem Clin Biochem* 1994;32:71-7.
284. Mbiti MJ, Orinda DA, Ojwang PJ. Reference intervals for some biochemical parameters in the aged Kenyan population. *East Afr Med J* 1994;71:84-7.
285. Franchini I, Bergamaschi E. [Biological indicators and reference values]. *G Ital Med Lav* 1994;16:5-9. [Italià]
286. Maranelli G. [Statistical aspects of the definition of reference values]. *G Ital Med Lav* 1994;16:31-6. [Italià]
287. Apostoli P. [Reference values for xenobiotics in biological matrixes: the state of the art]. *G Ital Med Lav* 1994;16:19-24. [Italià]
288. Andre M, Valsamis J. Reference values for androstanediol glucuronide. *Clin Chem* 1994;40:162. [Sense resum]
289. Yuregir GT, Aksoy K, Arpacı A, Unlukurt I, Tuli A. Studies on red cell glucose-6-phosphate dehydrogenase: evaluation of reference values. *Ann Clin Biochem* 1994;31:50-5.
290. Vernet M, Revenant MC, Charlier De Bressing C, Guillemain C, Pressac M, Renversez JC, Rymer JC. [Reference values of serum ferritin in adults]. *Ann Biol Clin* 1994;52:493-8. [Francès] [Sense resum]
291. Jimenez CV. Usefulness of reference limits and evaluation of significant differences. An example of the biological variation of serum rheumatoid factors. *Ann Biol Clin* 1994;52:529-33.
292. Stacewicz-Sapuntzakis M, Bowen PE, Mares-Perlman JA. Serum reference values for lutein and zeaxanthin using a rapid separation technique. *Ann N Y Acad Sci* 1993;691:207-9. [Sense resum]
293. Leung FY, Bradley C, Pellar TG. Reference intervals for blood lead and evaluation of zinc protoporphyrin as a screening test for lead toxicity. *Clin Biochem* 1993;26:491-6.
294. Vanhoe H, Versieck J, Vanballenberghe L, Dams R. Bismuth in human serum: reference interval and concentrations after intake of a therapeutic dose of colloidal bismuth subcitrate. *Clin Chim Acta* 1993;219:79-91.
295. Alstrom T, Grasbeck R, Lindblad B, Solberg HE, Winkel P, Viinikka L. Establishing reference values from adults: recommendation on procedures for the preparation of individuals, collection of blood, and handling and storage of specimens. Committee on Reference Values of the Scandinavian Society for Clinical Chemistry. *Scand J Clin Lab Invest* 1993;53:649-52.
296. Nagayama I, Yamamoto K, Saito K, Kuzuya T, Saito T. Subject-based reference values in thyroid function tests. *Endocr J* 1993;40:557-62.
297. Tholen DW. Reference values and participant means as targets in proficiency testing. *Arch Pathol Lab Med* 1993;117:885-9. [Revisió]

298. Boulat O, Janin B, Francioli P, Bachmann C. Plasma carnitines: reference values in an ambulatory population. *Eur J Clin Chem Clin Biochem* 1993;31:585-9.
299. Richardson-Jones A, Twedt D, Gibson M, Hellman R. Continuum analysis method applied to blood count reference intervals. *Ann Clin Lab Sci* 1993;23:340-9.
300. Solberg HE. A guide to IFCC recommendations on reference values. *J Int Fed Clin Chem* 1993;5:162-5. [Revisió]
301. Hoffmann GF, Seppel CK, Holmes B, Mitchell L, Christen HJ, Hanefeld F, Rating D, Nyhan WL. Quantitative organic acid analysis in cerebrospinal fluid and plasma: reference values in a pediatric population. *J Chromatogr* 1993;617:1-10.
302. Nuttall KL, Gordon WH, Ash KO. Whole-blood lead reference intervals for adults. *Clin Chem* 1993;39:1349-50. [Sense resum]
303. Perkins SL, Livesey JF, Belcher J. Reference intervals for 21 clinical chemistry analytes in arterial and venous umbilical cord blood. *Clin Chem* 1993;39:1041-4.
304. Dorner K, Schulze S. [Reference values for plasma amino acids in the course of pregnancy]. *Z Geburtshilfe Perinatol* 1993;197:141-3. [Alemany]
305. Siest G, Visvikis S, Zaiou M, Regis A, Jaid M, Steinmetz J, Herbeth B, Artur Y, Schiele F, Gueguen R, et al. The role of genetics in defining reference values and health status. *Methods Inf Med* 1993;32:255-9.
306. Blennow K, Fredman P, Wallin A, Gottfries CG, Karlsson I, Langstrom G, Skoog I, Svennerholm L, Wikkelso C. Protein analysis in cerebrospinal fluid. II. Reference values derived from healthy individuals 18-88 years of age. *Eur Neurol* 1993;33:129-33.
307. Djurhuus MS, Petersen PH, Rohold A, Vadstrup S, Uldall A. The basis for common reference intervals for serum potassium. *Ups J Med Sci* 1993;98:387-93.
308. Saitta M, Iavarone A, Cappello N, Bergami MR, Fiorucci GC, Aguzzi F. Reference values for immunoglobulin kappa and lambda light chains and the kappa/lambda ratio in children's serum. *Clin Chem* 1992;38:2454-7.
309. Serafini O, Battista V, Delprete E, Sanchini L, Gasperoni G, Galli F, Canestrari F. [Evaluation of an immunoenzymatic method for determining TSH and establishment of reference values]. *Boll Soc Ital Biol Sper* 1992;68:761-7. [Italià]
310. Tarallo P, Henny J, Gueguen R, Siest G. Reference limits of plasma fibrinogen. *Eur J Clin Chem Clin Biochem* 1992;30:745-51.
311. Lepage R, Whittom S, Bertrand S, Bahsali G, D'Amour P. Superiority of dynamic over static reference intervals for intact, midmolecule, and C-terminal parathyrin in evaluating calcemic disorders. *Clin Chem* 1992;38:2129-35.
312. Sugita O, Uchiyama K, Yamada T, Sato T, Okada M, Takeuchi K. Reference values of serum and urine creatinine, and of creatinine clearance by a new enzymatic method. *Ann Clin Biochem* 1992;29:523-8.
313. Fitzgibbon M, FitzGerald RJ, Tormey WP, O'Meara A, Kenny D. Reference values for urinary HMMA, HVA, noradrenaline, adrenaline, and dopamine excretion in children using random urine samples and HPLC with electrochemical detection. *Ann Clin Biochem* 1992;29:400-4.
314. Sasse EA. Determination of reference intervals in the clinical laboratory using the proposed guideline National Committee for Clinical Laboratory Standards C28-P. *Arch Pathol Lab Med* 1992;116:710-3. [Revisió]
315. Young DS. Determination and validation of reference intervals. *Arch Pathol Lab Med* 1992;116:704-9. [Revisió]

316. Ascione L, De Leva F, Cuomo S, Scherillo M, Caso P, Cioppa L, Mininni N. [Reference values for the echocardiographic calculation of left ventricular mass in normal children between the ages of 0 and 6]. *G Ital Cardiol* 1992;22:829-34. [Italià]
317. Bonadio WA, Stanco L, Bruce R, Barry D, Smith D. Reference values of normal cerebrospinal fluid composition in infants ages 0 to 8 weeks. *Pediatr Infect Dis J* 1992;11:589-91. [Sense resum]
318. Apostoli P. Criteria for the definition of reference values for toxic metals. *Sci Total Environ* 1992;120:23-37.
319. Grandjean P, Nielsen GD, Jorgensen PJ, Horder M. Reference intervals for trace elements in blood: significance of risk factors. *Scand J Clin Lab Invest* 1992;52:321-37.
320. Djurhuus MS, Rohold A, Vadstrup S, Petersen PH, Uldall A. Reference intervals based on hospitalized 'healthy' patients and medical students in relation to analytical bias for serum potassium. *Scand J Clin Lab Invest* 1992;52:305-12.
321. Marild S, Jodal U, Jonasson G, Mangelus L, Oden A, Persson NG. Reference values for renal concentrating capacity in children by the desmopressin test. *Pediatr Nephrol* 1992;6:254-7.
322. Liappis N, Schlebusch H, Niesen M. [Reference values for IgE concentration in serum of children. Method: ImmunoCAP-FEIA system]. *Monatsschr Kinderheilkd* 1992;140:300-2. [Alemany]
323. Pane F, Oriani G, Kuo KC, Gehrke CW, Salvatore F, Sacchetti L. Reference intervals for eight modified nucleosides in serum in a healthy population from Italy and the United States. *Clin Chem* 1992;38:671-7.
324. Whitehead RG. Dietary reference values. *Proc Nutr Soc* 1992;51:29-34. [Revisió] [Sense resum]
325. Wellington M, Mahomed K, Ngoro E. Reference values for glucose tolerance test in the urban Zimbabwean pregnant woman. *Cent Afr J Med* 1992;38:182-4.
326. Guarino A, Tarallo L, Greco L, Cesarano L, Guandalini S, Rubino A. Reference values of the steatocrit and its modifications in diarrheal diseases. *J Pediatr Gastroenterol Nutr* 1992;14:268-74.
327. Zunic G, Jelic-Ivanovic Z, Spasic S, Stojiljkovic A, Majkic-Singh N. Reference values for apolipoproteins A-I and B in healthy subjects, by age. *Clin Chem* 1992;38:566-9.
328. Herkner KR, Salzer H, Bock A, Muhl A, Tsaka T, Steger H, Pollak A, Schatten C, Popp W, Huber-Spitz V, et al. Pediatric and perinatal reference intervals for immunoglobulin light chains kappa and lambda. *Clin Chem* 1992;38:548-50.
329. Giannini SD, Deveriacki BE, De Gois JM, Diamant J, Forti N, Cardoso RH, Galluci SD, De Serro-Azul LG. [Prevalence of primary dyslipidemia in subjects with and without family history of coronary disease, based on the reference values of the "National Cholesterol Education Program" (NCEP)]. *Arq Bras Cardiol* 1992;58:281-7. [Portuguès]
330. Jung K, Pergande M, Schreiber G, Thierfelder W. Reference intervals for alpha 1-microglobulin in urine. *Clin Chim Acta* 1992;206:245-7. [Sense resum]
331. Schulze S, Dorner K. [Reference values for apolipoproteins and HDL-2/HDL-3 in childhood]. *Klin Padiatr* 1992;204:102-6. [Alemany]
332. Lozano JM, Duque OR, Buitrago T, Behaine S. Pulse oximetry reference values at high altitude. *Arch Dis Child* 1992;67:299-301.
333. Gibney MJ. Dietary reference values. *Nutr Rev* 1992;50:89. [Sense resum]
334. Kairisto V, Koskinen P, Mattila K, Puikkonen J, Virtanen A, Kantola I, Irjala K. Reference intervals for 24-h urinary normetanephrine, metanephrine, and 3-methoxy-4-hydroxymandelic acid in hypertensive patients. *Clin Chem* 1992;38:416-20.
335. Malvy DJ, Poveda JD, Debruyne M, Montagnon B, Burtschy B, Herbert C, Caces P, Houot O, Amedee-Manesme O. Laser immunonephelometry reference intervals for eight serum proteins in healthy children. *Clin Chem* 1992;38:394-9.

336. Porto G, Vicente C, Fraga J, da Silva BM, de Sousa M. Importance of establishing appropriate local reference values for the screening of hemochromatosis: a study of three different control populations and 136 hemochromatosis family members. Hemochromatosis Clinical and Research Group. *J Lab Clin Med* 1992;119:295-305.
337. Liappis N, Schlebusch H, Knapp M. [Reference values for the concentration of free triiodothyronine (FT3), triiodothyronine (TT3), free thyroxine (FT4), thyroxine (TT4), thyrotropin (TSH) and thyroxine-binding globulin (TBG) in umbilical cord blood. Method: luminescence-enhanced enzyme immunoassay]. *Klin Padiatr* 1992;204:34-6. [Alemany]
338. Wiedemann G, Simon T, Hildebrand U. [Establishment of reference values for fructosamine and HbA1c in childhood]. *Eur J Clin Chem Clin Biochem* 1992;30:33-8. [Alemany]
339. Jung K, Pergande M. Sex- and age-dependent reference values of alpha-1-microglobulin in urine. *Nephron* 1992;62:474-5. [Sense resum]
340. Lotter MG, Rabe WL, de Lange AH, Naude H, Otto AC, Meyer B, Heyns AD, Badenhorst PN. Reference values for red cell survival times. *J Nucl Med* 1991;32:2245-8.
341. Solberg HE, Stamm D. International Federation of Clinical Chemistry IFCC. IFCC recommendation--theory of reference values. Part 4. Control of analytical variation in the production, transfer and application of reference values. *Clin Chim Acta* 1991;202:S5-11. [Sense resum]
342. Fabre D, Horta H, Diez G, Escobar JP, Barrios F. Reference values of immunoglobulin E in umbilical cord sera. *J Investig Allergol Clin Immunol* 1991;1:335-9.
343. Carchon HA, Jaeken J, Jansen E, Eggermont E. Reference values for free gamma-aminobutyric acid determined by ion-exchange chromatography and fluorescence detection in the cerebrospinal fluid of children. *Clin Chim Acta* 1991;201:83-8. [Sense resum]
344. Harris EK, Wong ET, Shaw ST Jr. Statistical criteria for separate reference intervals: race and gender groups in creatine kinase. *Clin Chem* 1991;37:1580-2.
345. Solberg HE, Stamm D. International Federation of Clinical Chemistry, Scientific Division: approved recommendation on the theory of reference values. Part 4. Control of analytical variation in the production, transfer and application of reference values. *Eur J Clin Chem Clin Biochem* 1991;29:531-5. [Sense resum]
346. Pressac M, Morgant G, Farnier MA, Aymard P. Enzyme immunoassay of serum erythropoietin in healthy children: reference values. *Ann Clin Biochem* 1991;28:345-50.
347. Wilson DM, Liedtke RR. Modified enzyme-based colorimetric assay of urinary and plasma oxalate with improved sensitivity and no ascorbate interference: reference values and sample handling procedures. *Clin Chem* 1991;37:1229-35.
348. Noma A, Hata Y, Goto Y. Quantitation of serum apolipoprotein A-I, A-II, B, C-II, C-III and E in healthy Japanese by turbidimetric immunoassay: reference values, and age- and sex-related differences. *Clin Chim Acta* 1991;199:147-57.
349. Panaro A, Amati A, di Loreto M, Felle R, Ferrante M, Papadia AM, Porfido N, Gambatesa V, Dell'Osso A, Lucivero G. Lymphocyte subpopulations in pediatric age. Definition of reference values by flow cytometry. *Allergol Immunopathol* 1991;19:109-12.
350. Duca P. [Reference values and tolerance intervals]. *Med Lav* 1991;82:195-212. [Italià]
351. Piedras Ros J, Reyes Devesa S, Cordova Caballero MS, Chaves Rodriguez L. [Erythrocyte reference values in a Coulter S-Plus STKR, in healthy adults living at 2,240 meters above sea level]. *Rev Invest Clin* 1991;43:174-8. [Espanyol]
352. Berg B, Fex G, Tryding N, Nilsson JE, Ekman S. Reference intervals for retinol-binding protein in serum and urine. *Clin Chim Acta* 1991;197:149-52. [Sense resum]
353. van der Ploeg RJ, Fidler V, Oosterhuis HJ. Hand-held myometry: reference values. *J Neurol Neurosurg Psychiatry* 1991;54:244-7.

354. Liappis N, Schlebusch H, von Perjes M, Berg I. [Reference values for blood levels of free thyroxine, free triiodothyronine and thyroxine-binding globulins in euthyroid children. Method: luminescence-enhanced enzyme immunoassay]. *Klin Padiatr* 1991;203:113-5. [Alemany]
355. Lutz CT, Cornell SH, Goeken JA. Establishment of a reference interval for beta 2-microglobulin in cerebrospinal fluid with use of two commercial assays. *Clin Chem* 1991;37:104-7.
356. Klein G. [Results of multicenter determination of preliminary reference values for beta-N-acetylglucosaminidase in urine of adults]. *Wien Klin Wochenschr* 1991;189(Suppl):50-1. [Alemany]
357. Hubbuch A. [Results of multicenter determination of preliminary reference values for albumin in urine of children and adults]. *Wien Klin Wochenschr* 1991;189(Suppl.):48-9. [Alemany]
358. Szabolcs I, Ploenes C, Beyer M, Bernard W, Herrmann J. Reference intervals for serum thyrotropin: dependence on the population investigated. *Exp Clin Endocrinol* 1991;98:23-31.
359. Gelot MA. [Lysine in human biologic fluids and tissues: reference values, physiologic variations and interferences]. *Ann Biol Clin* 1991;49:53-6. [Francès]
360. Prunte C. [Reference values for quantitative image analysis of indocyanine green video-fluorescence angiography]. *Fortschr Ophthalmol* 1991;88:233-5. [Alemany]
361. Armstrong FM, Bradbury JE, Ellis SH, Owens DR, Rosen I, Sonksen P, Sundkvist G. A study of peripheral diabetic neuropathy. The application of age-related reference values. *Diabet Med* 1991;8(Spec No):S94-9.
362. Montalbetti N. Reference values in the nineties: from theory to practice? *Ann Ist Super Sanita* 1991;27:365-8. [Revisió]
363. Solberg HE, Stamm D. International Federation of Clinical Chemistry (IFCC) IFCC recommendation. The Theory of Reference Values. Part 4. Control of analytical variation in the production, transfer and application of reference values. *Ann Biol Clin* 1991;49:487-90. [Sense resum]
364. Ohba S, Kidouchi K, Nakamura C, Katoh T, Kobayashi M, Wada Y. Reference values of orotic acid, uracil and pseudouridine in urine. *Adv Exp Med Biol* 1991;309B:27-30. [Sense resum]
365. Beck T. Placental morphometry using a computer assisted measuring programme: reference values for normal pregnancies at term. *Arch Gynecol Obstet* 1991;249:135-47.
366. Anttila L, Koskinen P, Irjala K, Kaihola HL. Reference intervals for serum sex steroids and gonadotropins in regularly menstruating women. *Acta Obstet Gynecol Scand* 1991;70:475-81.
367. Flachaire E, Beney C, Berthier A, Salandre J, Quincy C, Renaud B. Determination of reference values for serotonin concentration in platelets of healthy newborns, children, adults, and elderly subjects by HPLC with electrochemical detection. *Clin Chem* 1990;36:2117-20.
368. Vicente C, Porto G, de Sousa M. Method for establishing serum ferritin reference values depending on sex and age. *J Lab Clin Med* 1990;116:779-84.
369. Pfammatter T, Marincek B, von Schulthess GK, Dudenhausen JW. [MR pelvimetric reference values]. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr* 1990;153:706-10. [Alemany]
370. Dale I. Plasma levels of the calcium-binding L1 leukocyte protein: standardization of blood collection and evaluation of reference intervals in healthy controls. *Scand J Clin Lab Invest* 1990;50:837-41.
371. Akiyama Y, Kazama M, Tahara C, Shimazu C, Otake J, Kamei K, Nakatake T, Sakurai N, Yasumuro Y, Suzuki S, et al. Reference values of hemostasis related factors of healthy Japanese adults. I: Circadian fluctuation. *Thromb Res* 1990;60:281-9.
372. Liappis N, Gobien N, Schlebusch H, Knapp M. [Age-specific and sex-specific reference values for the excretion of free amino acids in the 24-hour urine of children]. *Klin Padiatr* 1990;202:417-21. [Alemany]
373. Scott PH, Sandham S, Balmer SE, Wharton BA. Diet-related reference values for plasma amino acids in newborns measured by reversed-phase HPLC. *Clin Chem* 1990;36:1922-7.

374. Tarallo P, Henny J, Fournier B, Siest G. Plasma osteocalcin: biological variations and reference limits. *Scand J Clin Lab Invest* 1990;50:649-55.
375. Heeren TJ, Lagaay AM, von Beek WC, Rooymans HG, Hijmans W. Reference values for the Mini-Mental State Examination (MMSE) in octo- and nonagenarians. *J Am Geriatr Soc* 1990;38:1093-6.
376. Bester FC, Weich DJ, Barnard HC, Vermaak WJ. Biochemical reference values in elderly black subjects. *S Afr Med J* 1990;78:241-4.
377. Irjala K, Koskinen P, Icen A, Palosuo T. Reference intervals for immunoglobulins IgA, IgG and IgM in serum in adults and in children aged 6 months to 14 years. *Scand J Clin Lab Invest* 1990;50:573-7.
378. Chakar A, Mokni R, Suquet JP, Mahu JL, Bleiberg-Daniel F. Reference interval for transthyretin in children ages six to 60 months. *Clin Chem* 1990;36:1694. [Sense resum]
379. Honour JW. Steroid profile for urine: reference values. *Clin Chem* 1990;36(8 Pt 1):1528-9. [Sense resum]
380. Oosterhuis WP, Modderman TA, Pronk C. Reference values: Bhattacharya or the method proposed by the IFCC? *Ann Clin Biochem* 1990;27 (Pt 4):359-65.
381. Geven WB, Vogels-Mentink GM, Willems JL, de Boo T, Lemmens W, Monnens LA. Reference values of magnesium and potassium in mononuclear cells and erythrocytes of children. *Clin Chem* 1990;36:1323-7.
382. Masse J. Clinical chemistry reference intervals for healthy elderly subjects. *Am J Clin Nutr* 1990;51:1115-6. [Sense resum]
383. Van Hoof V, Geeraerts L, Van Meirvenne H, Van Mullem M, Chevigne R, Van Campenhout C, Lepoutre L. Reference values for amylase isoenzymes determined by agarose electrophoresis. *Clin Chem* 1990;36:1251-2. [Sense resum]
384. Takeoka T, Shinohara Y, Mori K, Furumi K. Solid-phase immunofluorometric assay for quantification of CSF immunoglobulins. Determination of normal reference values. *J Neurol Sci* 1990;96(2-3):229-40.
385. Liappis N, Gobien N, Schlebusch H. [Reference values of fasting serum levels of free amino acids in children]. *Klin Padiatr* 1990;202:161-7. [Alemany]
386. Belboul A, al-Khaja N, Bergman P, Roberts D, William-Olsson G. Blood cell filtrability: reference values and clinical applications. *Scand J Clin Lab Invest* 1990;50:297-302.
387. Kruesi MJ, Lenane MC, Hibbs ED, Major J. Normal controls and biological reference values in child psychiatry: defining normal. *J Am Acad Child Adolesc Psychiatry* 1990;29:449-52.
388. Duncanson GO, Worth HG. Determination of reference intervals for serum magnesium. *Clin Chem* 1990;36:756-8.
389. Atanasov NA. Graphic representation of reference intervals. *Clin Chem* 1990;36:701-2. [Sense resum]
390. Steinmetz J, Choukaife A, Visvikis S, Henny J, Siest G. Biological factors affecting concentrations of serum LpAI lipoprotein particles in serum, and determination of reference limits. *Clin Chem* 1990;36:677-80.
391. Whitley RJ. Reference values in pediatric medicine. *Mayo Clin Proc* 1990;65:431-5. [Sense resum]
392. Burritt MF, Slockbower JM, Forsman RW, Offord KP, Bergstralh EJ, Smithson WA. Pediatric reference intervals for 19 biologic variables in healthy children. *Mayo Clin Proc* 1990;65:329-36.
393. Liappis N, Schlebusch H. [Reference values of the ferritin concentration in the serum in children]. *Klin Padiatr* 1990;202:99-102. [Alemany]
394. Lindblad B, Alstrom T, Bo Hansen A, Grasbeck R, Hertz H, Holmberg C, Leskinen E, Moe PJ, Nyberg AP, Nanto V, et al. Recommendation for collection of venous blood from children, with special reference to production reference values. *Scand J Clin Lab Invest* 1990;50:99-104.

395. Beneteau-Burnat B, Baudin B, Morgant G, Baumann FC, Giboudeau J. Serum angiotensin-converting enzyme in healthy and sarcoidotic children: comparison with the reference interval for adults. *Clin Chem* 1990;36:344-6.
396. Kunkel S, Rudolf K, Buttner S, Wilms K. [Heterologous and homologous radioimmunoassays of human prolactin.. Reference values]. *Zentralbl Gynakol* 1990;112:629-32. [Alemany]
397. Henny J, Schiele F. [Age dependence, sex independence and reference values of serum fructosamine determined using a new colorimetry method]. *Wien Klin Wochenschr* 1990;180(Suppl.):48-52; discussion 57-9. [Alemany]
398. Rietdorf M, Hase I. [Specialty specific reference values in linen use in public health and social service as a basis for determining laundry capacities]. *Z Gesamte Hyg* 1990;36:35-7. [Alemany]
399. Knight JA, McClellan L, Staheli JK. Cerebrospinal fluid lipoperoxides quantified by liquid chromatography, and determination of reference values. *Clin Chem* 1990;36:139-42.
400. Ferner S, Koszmagk R, Lehmann A, Heilmann W. [Reference values of Na(+) and Cl(-) concentrations in adult sweat]. *Z Erkr Atmungsorgane* 1990;175:70-5. [Alemany]
401. Coni E, Falconieri P, Ferrante E, Semeraro P, Beccaloni E, Stacchini A, Caroli S. Reference values for essential and toxic elements in human milk. *Ann Ist Super Sanita* 1990;26:119-30.
402. Grasbeck R. Reference values, why and how. *Scand J Clin Lab Invest* 1990;201(Suppl.):45-53. [Revisió]
403. Lewenhaupt A, Ekman P, Eneroth P, Kallner A. Tumor markers in human prostatic carcinoma. An optimization of reference values. *Eur Urol* 1990;17:7-11.
404. Chen ZQ, Yu JH, Cao SH. Reference values of indicators for WHO neurobehavioral core test battery. *Chin Med J* 1990;103:61-5.
405. Kanabrocki EL, Sothorn RB, Scheving LE, Vesely DL, Tsai TH, Shelstad J, Cournoyer C, Greco J, Mermall H, Ferlin H, et al. Reference values for circadian rhythms of 98 variables in clinically healthy men in the fifth decade of life. *Chronobiol Int* 1990;7(5-6):445-61.
406. Siggaard-Andersen O, Wimberley PD, Fogh-Andersen N, Gothgen IH. Arterial oxygen status determined with routine pH/blood gas equipment and multi-wavelength hemoximetry: reference values, precision, and accuracy. *Scand J Clin Lab Invest* 1990;203(Suppl.):57-66.
407. Farin G. [Physical differentiation of methods of electrocoagulation with reference to specific limits of performance]. *Biomed Tech* 1990;35(Suppl 3):198-9. [Alemany] [Sense resum]
408. Berger J. Computer evaluation of reference limits. *Folia Haematol Int Mag Klin Morphol Blutforsch* 1990;117:475-8.
409. Hyltoft Petersen P, Gowans EM, Blaabjerg O, Horder M. Analytical goals for the estimation of non-Gaussian reference intervals. *Scand J Clin Lab Invest* 1989;49:727-37.
410. Weykamp CW, Penders TJ, Schmidt NA, Borburgh AJ, van de Calseyde JF, Wolthers BJ. Steroid profile for urine: reference values. *Clin Chem* 1989;35:2281-4.
411. Garry PJ, Hunt WC, VanderJagt DJ, Rhyne RL. Clinical chemistry reference intervals for healthy elderly subjects. *Am J Clin Nutr* 1989;50(5 Suppl):1219-30; discussion 1231-5.
412. Jarvisalo J, Maatela J, Maki J, Marniemi J, Reunanen A. Health-based reference values of the Mini-Finland Health Survey: 1. Serum gamma-glutamyltransferase, aspartate aminotransferase and alkaline phosphatase. *Scand J Clin Lab Invest* 1989;49:623-32.
413. Nicolau GY, Haus E. Chronobiologic reference values in clinical chemistry. *Endocrinologie* 1989;27:197-230.
414. Loria A, Salas R. [Patient data for evaluating reference values]. *Rev Invest Clin* 1989;41:381-5. [Espanyol]

415. Scott PH. Enzyme immunoassay of lactoferrin in newborn term infants: reference values and influence of diet. *Ann Clin Biochem* 1989;26:407-11.
416. Delanghe J, De Slypere JP, De Buyzere M, Robbrecht J, Wieme R, Vermeulen A. Normal reference values for creatine, creatinine, and carnitine are lower in vegetarians. *Clin Chem* 1989;35:1802-3. [Sense resum]
417. Lott JA, Warren P. Estimation of reference intervals for total protein in cerebrospinal fluid. *Clin Chem* 1989;35:1766-70.
418. Gerrits GP, Trijbels FJ, Monnens LA, Gabreels FJ, De Abreu RA, Theeuwes AG, van Raay-Selten B. Reference values for amino acids in cerebrospinal fluid of children determined using ion-exchange chromatography with fluorimetric detection. *Clin Chim Acta* 1989;182:271-80.
419. Weinmann G, Menzel K. [Immunologic findings in the newborn infant. II. Studies of humoral immunity, reference values of immunologic studies in the newborn infant]. *Kinderarztl Prax* 1989;57:327-33. [Alemany]
420. Lindstedt G, Lundberg PA, Johansson PM, Eggertsen R, Ellertz G, Nystrom E. High prevalence of atrophic gastritis in the elderly: implications for health-associated reference limits for cobalamin in serum. *Clin Chem* 1989;35:1557-8. [Sense resum]
421. Gimsing P, Nexø E. Cobalamin-binding capacity of haptocorrin and transcobalamin: age-correlated reference intervals and values from patients. *Clin Chem* 1989;35:1447-51.
422. Chomon B, Vazquez L, Castro D. [Reference intervals of hematologic parameters in the elderly]. *Sangre* 1989;34:229-33. [Espanyol]
423. Plouin PF, Chatellier G, Guyene TT, Vincent N, Corvol P. [Recent advances in the clinical study of the renin system. Reference values and conditions of validity]. *Presse Med* 1989;18:917-21. [Francès]
424. Larsson L, Ohman S, Nelson N, Finnstrom O. Neonatal reference values for ionized calcium. Some comments on criteria in selection of reference population. *Scand J Clin Lab Invest* 1989;49:301-2. [Sense resum]
425. Feldmann D, Flandrois C, Jardel A, Phan TM, Aymard P. Circadian variations and reference intervals for some enzymes in urine of healthy children. *Clin Chem* 1989;35:864-7.
426. Hasselot N, Ott D, Blanchot M. [Determination of some usual and frequent values in clinical biochemistry in a Polynesian population sample. Comparison with French reference values]. *Med Trop* 1989;49:179-82. [Francès]
427. Gimpel JA, van Rijn HJ, Putters J. Reference interval for the bilirubin excess in cerebrospinal fluid by derivative spectrophotometry. *J Clin Chem Clin Biochem* 1989;27:217-9.
428. Lambert WE, Vanneste L, De Leenheer AP. Serum phylloquinone concentrations in a normal population of children as compared with the reference interval in adults. *Clin Chem* 1989;35:671. [Sense resum]
429. López JB. Reference values terminology. *Med Lab Sci* 1989;46:161. [Sense resum]
430. Junge W, Troge B, Klein G, Poppe W, Gerber M. Evaluation of a new assay for pancreatic amylase: performance characteristics and estimation of reference intervals. *Clin Biochem* 1989;22:109-14.
431. Gerver WJ, Drayer NM, Schaafsma W. Reference values of anthropometric measurements in Dutch children. The Oosterwolde Study. *Acta Paediatr Scand* 1989;78:307-13.
432. Spichiger UE, Vonderschmitt DJ. A self-consistent set of reference values for 23 clinical chemical analytes. *Clin Chem* 1989;35:448-52.
433. Morris CD, Dickson D. Haematological reference values in adult blacks at sea level. *S Afr Med J* 1989;75:35-6. [Sense resum]

434. Panteghini M, Pagani F. Reference intervals for two bone-derived enzyme activities in serum: bone isoenzyme of alkaline phosphatase (ALP) and tartrate-resistant acid phosphatase (TR-ACP). *Clin Chem* 1989;35:180-1. [Sense resum]
435. Nordin BE, Need AG, Hartley TF, Philcox JC, Wilcox M, Thomas DW. Improved method for calculating calcium fractions in plasma: reference values and effect of menopause. *Clin Chem* 1989;35:14-7.
436. Wehner HD, Schieffer MC. [Reference values of n-propanol elimination]. *Beitr Gerichtl Med* 1989;47:385-9. [Alemany]
437. Lawrence C, Fryer JG, Karlberg P, Niklasson A, Ericson A. Modelling of reference values for size at birth. *Acta Paediatr Scand* 1989;350(Suppl.):55-69.
438. Sandhagen B. Analysis of haemorheological variables--methodology and reference values. *Ups J Med Sci* 1989;94:81-7.
439. Solberg HE, Grasbeck R. Reference values. *Adv Clin Chem* 1989;27:1-79. [Revisió] [Sense resum]
440. Guillemín C, Plomteux G, Dezier JF, Paris M, Pressac M, Revenant MC, Vernet M. [Reference values of erythrocyte ferritin in children and adults]. *Ann Biol Clin* 1989;47:203-6. [Revisió] [Francès]
441. Krantz S, Lober M, Lober S. [The determination of the reference range and the reference limits of the fructosamine test]. *Z Med Lab Diagn* 1989;30:395-6. [Alemany] [Sense resum]
442. Voss P, Linneke P, Haller G, Jeske W. [Reference values for progesterone, beta-HCG, estriol and human placental lactogen in the 10th-16th weeks of pregnancy]. *Zentralbl Gynakol* 1989;111(13):897-903. [Alemany]
443. De Schepper J, Derde MP, Goubert P, Gorus F. Reference values for fructosamine concentrations in children's sera: influence of protein concentration, age, and sex. *Clin Chem* 1988;34:2444-7.
444. Lepage R, D'Amour P, Boucher A, Hamel L, Demontigny C, Labelle F. Clinical performance of a parathyrin immunoassay with dynamically determined reference values. *Clin Chem* 1988;34:2439-43.
445. Hurst PL, Langford NJ, Shue J. Reference intervals for plasma alanine aminopeptidase in the Hitachi 737 analyzer, derived from patients with normal biochemical liver profiles. *Clin Chem* 1988;34:2583-4. [Sense resum]
446. Bingham SA, Williams R, Cole TJ, Price CP, Cummings JH. Reference values for analytes of 24-h urine collections known to be complete. *Ann Clin Biochem* 1988;25:610-9.
447. Parvy PR, Bardet JI, Rabier DM, Kamoun PP. Age-related reference values for free amino acids in first morning urine specimens. *Clin Chem* 1988;34:2092-5.
448. Bais R, Conyers RA, Rofe AM, Tormet RI, Geary TD. Creatine kinase reference intervals determined from a multi-centre data pool. *Pathology* 1988;20:367-72.
449. Yatscoff RW, Dube WJ. Reference interval for basal calcitonin and calcitonin concentrations after pentagastrin infusion. *Clin Chem* 1988;34:1931-2. [Sense resum]
450. Sherry B, Jack RM, Weber A, Smith AL. Reference interval for prealbumin for children two to 36 months old. *Clin Chem* 1988;34:1878-80.
451. Solberg HE, PetitClerc C. International Federation of Clinical Chemistry (IFCC), Scientific Committee, Clinical Section, Expert Panel on Theory of Reference Values. Approved recommendation (1988) on the theory of reference values. Part 3. Preparation of individuals and collection of specimens for the production of reference values. *J Clin Chem Clin Biochem* 1988;26:593-8. [Sense resum]
452. Liappis N, Schlebusch H. [Reference values for the concentrations of triiodothyronine, thyroxine and thyrotropin in the blood serum of euthyroid children. Method: luminescence-enhanced enzyme immunoassay]. *Klin Padiatr* 1988;200:410-3. [Alemany]

453. Campbell AJ, Borrie MJ. Reference values for upper arm anthropometric measurements for a New Zealand community sample of subjects aged 70 years and over. *Hum Biol* 1988;60:587-96. [Sense resum]
454. Nilsson-Ehle H, Jagenburg R, Landahl S, Svanborg A, Westin J. Haematological abnormalities in a 75-year-old population. Consequences for health-related reference intervals. *Eur J Haematol* 1988;41:136-46.
455. Lockitch G, Halstead AC, Wadsworth L, Quigley G, Reston L, Jacobson B. Age- and sex-specific pediatric reference intervals and correlations for zinc, copper, selenium, iron, vitamins A and E, and related proteins. *Clin Chem* 1988;34:1625-8.
456. Lockitch G, Halstead AC, Albersheim S, MacCallum C, Quigley G. Age- and sex-specific pediatric reference intervals for biochemistry analytes as measured with the Ektachem-700 analyzer. *Clin Chem* 1988;34:1622-5.
457. Lockitch G, Halstead AC, Quigley G, MacCallum C. Age- and sex-specific pediatric reference intervals: study design and methods illustrated by measurement of serum proteins with the Behring LN Nephelometer. *Clin Chem* 1988;34:1618-21.
458. Gerrits GP, Haagen AA, De Abreu RA, Monnens LA, Gabreels FJ, Trijbels FJ, Theeuwes AL, van Baal JM. Reference values for nucleosides and nucleobases in cerebrospinal fluid of children. *Clin Chem* 1988;34:1439-42.
459. Sanderink GJ, Artur Y, Schiele F, Gueguen R, Siest G. Alanine aminopeptidase in serum: biological variations and reference limits. *Clin Chem* 1988;34:1422-6.
460. Evans GO. Reference intervals for angiotensin-converting enzymes. *Clin Chem* 1988;34:1523-4. [Sense resum]
461. Leroux ML, Desjardins PR. Reference intervals for serum magnesium determined with a DuPont aca. *Clin Chem* 1988;34:1363. [Sense resum]
462. Knight JA, Smith SE, Kinder VE, Pieper RK. Urinary lipoperoxides quantified by liquid chromatography, and determination of reference values for adults. *Clin Chem* 1988;34:1107-10.
463. Thompson RS, Trudinger BJ, Cook CM, Giles WB. Umbilical artery velocity waveforms: normal reference values for A/B ratio and Pourcelot ratio. *Br J Obstet Gynaecol* 1988;95:589-91.
464. Fernandez de Castro M, Latorre P, Tafalla C, Vilorio A, Fernandez P, Larrocha C, Jimenez MC. [Reference values of basic hematologic parameters. An adult population]. *Sangre* 1988;33:188-95. [Espanyol] [Sense resum]
465. Seamonds B. Reference intervals for prostate-specific antigen. *Clin Chem* 1988;34:1366-7. [Sense resum]
466. Nasrat AA. Reference values for 75 g oral glucose tolerance test in pregnancy. *Br Med J (Clin Res Ed)* 1988;296:1330. [Sense resum]
467. Wandrup J, Kroner J, Pryds O, Kastrup KW. Age-related reference values for ionized calcium in the first week of life in premature and full-term neonates. *Scand J Clin Lab Invest* 1988;48:255-60.
468. Kuwana T, Sugita O, Yakata M. Reference limits of bone and liver alkaline phosphatase isoenzymes in the serum of healthy subjects according to age and sex as determined by wheat germ lectin affinity electrophoresis. *Clin Chim Acta* 1988;173:273-80.
469. [No authors listed] Reference values for 75 g oral glucose tolerance test in pregnancy. *Br Med J (Clin Res Ed)* 1988;296:1125-6. [Sense resum]
470. Rust RS Jr, Dodson WE, Trotter JL. Cerebrospinal fluid IgG in childhood: the establishment of reference values. *Ann Neurol* 1988;23:406-10.
471. Schiele F, Artur Y, Floc'h AY, Siest G. Total, tartrate-resistant, and tartrate-inhibited acid phosphatases in serum: biological variations and reference limits. *Clin Chem* 1988;34:685-90.

472. Watts GF, Morris RW, Khan K, Polak A. Urinary albumin excretion in healthy adult subjects: reference values and some factors affecting their interpretation. *Clin Chim Acta* 1988;172:191-8.
473. Hatem M, Anthony F, Hogston P, Rowe DJ, Dennis KJ. Reference values for 75 g oral glucose tolerance test in pregnancy. *Br Med J (Clin Res Ed)* 1988;296:676-8.
474. Iyengar V, Woittiez J. Trace elements in human clinical specimens: evaluation of literature data to identify reference values. *Clin Chem* 1988;34:474-81. [Revisió]
475. Maguire GA, Price CP. Reference intervals for the kinetic angiotensin converting enzyme assay. *Clin Chem* 1988;34:464. [Sense resum]
476. Grasbeck R. Reference values: relevant controls for clinical decision-making. *Eur J Haematol* 1988;40:1-6. [Sense resum]
477. Liappis N. [Reference values of beta 2 microglobulin concentrations in the serum of children]. *Klin Padiatr* 1988;200:67-9. [Alemany]
478. Nilsson-Ehle H, Jagenburg R, Landahl S, Svanborg A, Westin J. Haematological abnormalities and reference intervals in the elderly. A cross-sectional comparative study of three urban Swedish population samples aged 70, 75 and 81 years. *Acta Med Scand* 1988;224:595-604.
479. Lewenhaupt A, Ekman P, Eneroth P, Kallner A. On reference values for tumor markers in prostatic carcinoma. *Scand J Urol Nephrol* 1988;110(Suppl.):119-23.
480. Leipnitz G, Miyashita C, Heiden M, von Blohn G, Kohler M, Wenzel E. Reference values and variability of plasminogen in healthy blood donors and its relation to parameters of the fibrinolytic system. *Haemostasis* 1988;18(Suppl.) 1:61-8.
481. Martini E, Roquin H, Gastal C, Doinel C. [Reduction of circulating lymphocytes after giving blood. Effects on the establishment of reference values for CD4 and CD8 lymphocytes]. *Ann Biol Clin* 1988;46:327-8. [Francès] [Sense resum]
482. Nystrom E, Bengtsson C, Lindstedt G, Lapidus L, Lindquist O, Waldenstrom J. Serum gamma-glutamyltransferase in a Swedish female population. Age-related reference intervals; morbidity and prognosis in cases with raised catalytic concentration. *Acta Med Scand* 1988;224:79-84.
483. Ekman P, Lewenhaupt A, Eneroth P, Kallner A. Aspects on reference values for tumor markers in human prostatic carcinoma. *Am J Clin Oncol* 1988;11 Suppl 2:S80-2.
484. Beukeveld GJ, Wolthers BG, van Saene JJ, de Haan TH, de Ruyter-Buitenhuys LW, van Saene RH. Patterns of porphyrin excretion in feces as determined by liquid chromatography; reference values and the effect of flora suppression. *Clin Chem* 1987;33:2164-70.
485. Knight JA, Smith SE, Kinder VE, Anstall HB. Reference intervals for plasma lipoperoxides: age-, sex-, and specimen-related variations. *Clin Chem* 1987;33:2289-91.
486. Roulston JE. Different reference intervals for the kinetic assay of angiotensin-converting enzyme. *Clin Chem* 1987;33:2324-5. [Sense resum]
487. Duffau G, Leon A, Farias P, Urcelay G. [Fecal alpha 1-antitrypsin in infants. Reference values and in acute diarrheal syndrome]. *Rev Chil Pediatr* 1987;58:442-5. [Espanyol] [Sense resum]
488. Plomteux G, Charlier C, Albert A, Farnier M, Pressac M, Vernet M, Paris M, Dellamonica C, Dezier JF. [Reference values for serum transferrin in newborn infants and children]. *Ann Pediatr* 1987;34:649-52. [Francès] [Sense resum]
489. Liappis N, Starke A. [Reference values for the concentration of free thyroxine and free triiodothyronine in the serum of euthyroid children]. *Klin Padiatr* 1987;199:366-9. [Alemany]
490. Costongs GM, Hemrika MH, Engels LG, Bos LP, Bas BM, Flendrig JA, Janson PC. Faecal lysozyme: determination, reference intervals and some data in gastro-intestinal disease. *Clin Chim Acta* 1987;167:125-34.

491. Swaanenburg JC, Rutten WP, Holdrinet AC, van Strik R. The determination of reference values for hematologic parameters using results obtained from patient populations. *Am J Clin Pathol* 1987;88:182-91.
492. Gallacher RE, Browning MC, Fraser CG, Wilkinson SP, MacLennan WJ. A method for simultaneously estimating plasma, erythrocyte, and leukocyte sodium, potassium, and magnesium: reference values and considerations from biological variation data. *Clin Chem* 1987;33:1326-30.
493. BATTERY JE. Why are there different reference intervals for the kinetic angiotensin-converting enzyme assay? *Clin Chem* 1987;33:1491-2. [Sense resum]
494. Tikly M, Blumsohn D, Solomons HD, Govender Y, Atkinson PM. Normal haematological reference values in the adult black population of the Witwatersrand. *S Afr Med J* 1987;72:135-6.
495. Soliman SA, Abdel-Hay MH, Tayeb OS, Sulaiman MI. Application of transformation algorithm and nonparametric calculation in determining the reference intervals of some urine constituents and characteristics. *Clin Chim Acta* 1987;166:9-16.
496. Dionigi R, Berti Riboli E, Bertoglio S, Bonoldi AP, Braga M, Brandi LS, Coccaglio R, Corazza GR, Craus W, Cristallo M, et al. [Evaluation of the delayed hypersensitivity response by the Multitest-IMC. Multicenter study for the definition of reference values in a sample population]. *Minerva Med* 1987;78:721-31. [Italia]
497. Maturen A, Francoeur TA, Wynn AL, Reid DL, Anderson CP. Clinical performance of the Abbott "Vision System": reference intervals and specimen type comparisons. *Clin Chem* 1987;33:1078-9. [Sense resum]
498. Grisler R, Giavardi C, Scaricabarozzi R. [Reference values for sialic acid in the urine]. *Quad Sclavo Diagn* 1987;23:228-31. [Italia]
499. Georgieff M. [Intravenous feeding--possibilities and limits in the use of glucose and xylitol following trauma and infection, with special reference to liver metabolism]. *Infusionsther Klin Ernahr* 1987;14:93-7. [Alemany]
500. Solberg HE. International Federation of Clinical Chemistry (IFCC), Scientific Committee, Clinical Section, Expert Panel on Theory of Reference Values, and International Committee for Standardization in Haematology (ICSH), Standing Committee on Reference Values. Approved Recommendation (1986) on the theory of reference values. Part 1. The concept of reference values. *J Clin Chem Clin Biochem* 1987;25:337-42. [Sense resum]
501. Alstrom T, Dahl M, Grasbeck R, Hagenfeldt L, Hertz H, Hjelm M, Jarvenpaa AL, Kantero R, Larsson A, Leskinen EE, et al. Recommendation for collection of skin puncture blood from children, with special reference to production of reference values. *Scand J Clin Lab Invest* 1987;47:199-205.
502. Chiecchio A, Bo A, Migliardi M. Analysis of distributions and determination of reference limits in short series of data. Application to RIA. *J Nucl Med Allied Sci* 1987;31:195-200. [Sense resum]
503. Kallner A, Kallner G, Lundell G, Sjoberg HE. Highly sensitive assays of serum-thyrotropin in the diagnosis of hyperthyroidism: assessment of performance and reference values. *Scand J Clin Lab Invest* 1987;47:157-64.
504. Nelson N, Finnstrom O, Larsson L. Neonatal reference values for ionized calcium, phosphate and magnesium. Selection of reference population by optimality criteria. *Scand J Clin Lab Invest* 1987;47:111-7.
505. Kreuzfelder E, Shen G, Rodeck U, Konig E, Luboldt W, Keinecke HO, Brockmeyer N, Scheiermann N. Relative and absolute numbers of human lymphocyte subpopulations. A comparison of immunofluorescence microscopy and flow cytometric methodologies with special reference to precision and reference values. *J Immunol Methods* 1987;97:251-8.
506. Garry PJ. More on reference intervals for some vitamins. *Clin Chem* 1987;33:441-3. [Sense resum]
507. Zlotkin SH, Casselman CW. Percentile estimates of reference values for total protein and albumin in sera of premature infants (less than 37 weeks of gestation). *Clin Chem* 1987;33:411-3.

508. Franco G, Fidanza L, Casale G. Biological monitoring and chronobiology: preliminary time-quantified reference values of urinary zinc. *Chronobiologia* 1987;14:49-52. [Sense resum]
509. Gelot MA. [Amino acids in cerebrospinal fluid. Reference values]. *Ann Biol Clin* 1987;45:15-9. [Francès]
510. Rezuke WN, Knight JA, Sunderman FW Jr. Reference values for nickel concentrations in human tissues and bile. *Am J Ind Med* 1987;11:419-26.
511. Ohrloff C, Spitznas M. Reference values for normal corneal transfer coefficient (Kc) as tested with the Fluorotron-Master. *Graefes Arch Clin Exp Ophthalmol* 1987;225:84. [Sense resum]
512. Plomteux G, Charlier C, Albert A, Farnier M, Pressac M, Vernet M, Paris M, Dellamonica C, Dezier JF. [Reference values of serum transferrin in newborn infants, children and adults]. *Ann Biol Clin* 1987;45:622-9. [Francès]
513. Bergmann S, Heinrich JJ, Gehrlich S, Hanisch J, Tappert C, Beutner M, Jaross W. [Apolipoprotein concentrations in cord blood in healthy neonates--comparison with adult reference values]. *Z Med Lab Diagn* 1987;28:385-92. [Alemany] [Sense resum]
514. Schmitz JE, Lindner KH, Grunert A. [Reference values in the detection of disorders of homeostasis in shock]. *Klin Anasthesiol Intensivther* 1987;33:226-40. [Revisió] [Alemany] [Sense resum]
515. Lockitch G, McCallum C, Halstead C, Pendray M. Reference intervals for eight plasma proteins in preterm and term newborns by laser nephelometry. *Clin Chem* 1986;32:2106. [Sense resum]
516. Ng RH, Altaffer M, Statland BE. Reference intervals for nine tests on the RA-1000. *Clin Chem* 1986;32:1984. [Sense resum]
517. Osifo BO, Lukanmbi FA, Bolodeoku JO. Reference values for serum folate, erythrocyte folate and serum cobalamin in Nigerian adolescents. *Trop Geogr Med* 1986;38:259-64.
518. Herbeth B, Zittoun J, Miravet L, Bourgeay-Causse M, Carre-Guery G, Delacoux E, Le Devehat C, Lemoine A, Mareschi JP, Martin J, et al. Reference intervals for vitamins B1, B2, E, D, retinol, beta-carotene, and folate in blood: usefulness of dietary selection criteria. *Clin Chem* 1986;32:1756-9.
519. Tietz NW, Shuey DF. Reference intervals for alkaline phosphatase activity determined by the IFCC and AACC reference methods. *Clin Chem* 1986;32:1593-4. [Sense resum]
520. Premel-Cabic A, Turcant A, Allain P. Normal reference intervals for free catecholamines and their acid metabolites in 24-h urines from children, as determined by liquid chromatography with amperometric detection. *Clin Chem* 1986;32:1585-7.
521. Jain S. Reference interval for urinary glucose: effect of age. *Clin Chem* 1986;32:1426. [Sense resum]
522. Fenoil R, Weisz G, Alfieri R. Reference values for CK-MB isoenzyme compared with acute myocardial infarction. *Clin Chem* 1986;32:1238. [Sense resum]
523. Marima-Matarira HT. Reference values of serum protein fractions in Zimbabwe by electrophoresis. *Cent Afr J Med* 1986;32:116-8. [Sense resum]
524. Lakhani SA, Jansen AA, Gemert W. Some anthropometric values of Indian infants from Nairobi compared with infants from India, United Kingdom and the Harvard reference values. *East Afr Med J* 1986;63:304-11. [Sense resum]
525. Compton PJ, Stuart MC, Lazarus L. Error in laboratory reference limits as shown in a collaborative quality-assurance program. *Clin Chem* 1986;32:845-9.
526. Jain S. Reference interval for urinary glucose in elderly subjects. *Clin Chem* 1986;32:711-2. [Sense resum]
527. de Kieviet W, Slaats EH, Abeling NG. Pediatric reference values for calcium, magnesium and inorganic phosphorus in serum obtained from Bhattacharya plots for data from unselected patients. *J Clin Chem Clin Biochem* 1986;24:233-42.

528. Lorentz K, Weiss T, Kraas E. Sialic acid in human serum and cerebrospinal fluid. Comparison of methods and reference values. *J Clin Chem Clin Biochem* 1986;24:189-98.
529. Sinton TJ, Cowley DM, Bryant SJ. Reference intervals for calcium, phosphate, and alkaline phosphatase as derived on the basis of multichannel-analyzer profiles. *Clin Chem* 1986;32(1 Pt 1):76-9.
530. Hermosa V, Mazo E, Carril J, Cordovilla JJ, Luceno E, Zubizarreta A. [Reference values of analytical parameters used in the diagnosis of iron deficiency]. *Sangre* 1986;31:534-44. [Espanyol] [Sense resum]
531. Herbeth B, Henny J, Siest G. [Determination of reference values: a priori or a posteriori selection]? *Ann Biol Clin* 1986;44:411-2. [Francès] [Sense resum]
532. Bouman AA, Platenkamp AJ, Posma FD. Determination of aluminium in human tissues by flameless atomic absorption spectroscopy and comparison of reference values. *Ann Clin Biochem* 1986;23 (Pt 1):97-101.
533. Marima-Matarira HT. Reference values for urea, creatinine and electrolytes in Zimbabwean females. *Cent Afr J Med* 1986;32:21-2. [Sense resum]
534. Orts-Costa J, Juan-Pereira L, Fuentes-Arderiu J. Serum adenosine deaminase reference interval. *Ann Biol Clin* 1986;44:63. [Sense resum]
535. Grandjean P. Reference intervals for toxic metals: problems and prospects. Abraham J. Gitlitz memorial lecture. *Ann Clin Lab Sci* 1986;16:67-74. [Revisió]
536. Solberg HE. Statistical treatment of reference values in laboratory medicine: testing the goodness-of-fit of an observed distribution to the Gaussian distribution. *Scand J Clin Lab Invest* 1986;184(Suppl):125-32.
537. Canfell PC, Binder SR, Khayam-Bashi H. Pediatric reference intervals for normetanephrine/metanephrine. *Clin Chem* 1986;32:222-3. [Sense resum]
538. Nicholson GA, McLeod JG, Morgan G, Meerkin M, Cowan J, Bretag A, Graham D, Hill G, Robertson E, Sheffield L. Variable distributions of serum creatine kinase reference values. Relationship to exercise activity. *J Neurol Sci* 1985;71:233-45.
539. Nicholson GA, Morgan G, Meerkin M, Strauss E, McLeod JG. The creatine kinase reference interval. An assessment of intra- and inter-individual variation. *J Neurol Sci* 1985;71:225-31.
540. Juan-Pereira L, Andres-Moreno R, Fuentes-Arderiu J. Reference interval for serum alpha-amylase determined with p-nitrophenyl-alpha-D-maltoheptaoside as a substrate. *J Clin Chem Clin Biochem* 1985;23:861-3.
541. Shultz EK, Willard KE, Rich SS, Connelly DP, Critchfield GC. Improved reference-interval estimation. *Clin Chem* 1985;31:1974-8.
542. Marima-Matarira HT. Reference values for urea, creatinine and electrolytes in Zimbabwean males. *Cent Afr J Med* 1985;31:235-7. [Sense resum]
543. Baadenhuijsen H, Smit JC. Indirect estimation of clinical chemical reference intervals from total hospital patient data: application of a modified Bhattacharya procedure. *J Clin Chem Clin Biochem* 1985;23:829-39.
544. Neumeier D, Pauls R, Wifling I, Fateh-Moghadam A, Knedel M. [Immunophenotyping of lymphocyte subpopulations using monoclonal antibodies in an analytical flow cytometer: evaluation of the method and reference values]. *J Clin Chem Clin Biochem* 1985;23:765-75. [Alemany]
545. Sagniez M, Krempf M, Le Bastard N. [Correlation between the assay of glycosylated hemoglobin A1c and fructosamine in diabetic patients submitted to a systematic control visit. Establishment of reference values]. *Pathol Biol* 1985;33:914-6. [Francès]
546. Senan C, Rymer JC, Rosa J. [Assay of glycosylated hemoglobin A1c in children. Determination of reference values]. *Presse Med* 1985;14:1885. [Francès] [Sense resum]

547. Bandi ZL, Schoen I, Bee DE. Immunochemical methods for measurement of transferrin in serum: effects of analytical errors and inappropriate reference intervals on diagnostic utility. *Clin Chem* 1985;31:1601-5.
548. Cranendonk E, van Gennip AH, Abeling NG, Behrendt H, Hart AA. Reference values for automated cytochemical differential count of leukocytes in children 0-16 years old: comparison with manually obtained counts from Wright-stained smears. *J Clin Chem Clin Biochem* 1985;23:663-7.
549. Uhlhaas S, Olek K. [Concentrations of free and bound 4-aminobutyric acid in human serum: reference values]. *J Clin Chem Clin Biochem* 1985;23:525-8. [Alemany]
550. Gallo C, Turco MC, Sanzi G, La Rocca S, Sacchetti L, Salvatore F. Application of a statistical approach to determining reference intervals in clinical chemistry. *Clin Chem* 1985;31:1367-70.
551. Christenson RH, Dent GA, Tuszyński A. Two radioassays for serum vitamin B12 and folate determination compared in a reference interval study. *Clin Chem* 1985;31:1358-60.
552. Howe A, Thompson D, Wright V. Reference values for metacarpophalangeal joint stiffness in normals. *Ann Rheum Dis* 1985;44:469-76.
553. Melzi d'Eril GV, Pollini E, Moratti R, Sprovieri G, Tocchini M, Galeazzi M. Proposed reference values for fecal chymotrypsin as measured photometrically. *Clin Chem* 1985;31:1088-9. [Sense resum]
554. Cuvelier I, Steinmetz J, Mikstacki T, Siest G. Variations in total phospholipids and high-density lipoprotein phospholipids in plasma from a general population: reference intervals and influence of xenobiotics. *Clin Chem* 1985;31:763-6.
555. Bhattacharya RD, Patel TS, Pandya CB. Reference values of urinary electrolyte excretion in respect of a vegetarian diet. A chronobiological study. *Panminerva Med* 1985;27:83-7. [Sense resum]
556. Trundle D, Susko K. Reference interval for thyroxin as measured by fluorescence polarization immunoassay. *Clin Chem* 1985;31:650-1. [Sense resum]
557. Lepage L, Schiele F, Gueguen R, Siest G. Total cholinesterase in plasma: biological variations and reference limits. *Clin Chem* 1985;31:546-50.
558. Person B, Leglise D, Bardou LG. [On the need for establishing one's own reference values of apolipoproteins A1 and B in the detection of dyslipoproteinemia]. *Presse Med* 1985;14:642. [Francès] [Sense resum]
559. Behr W, Schlimok G, Firchau V, Paul HA. Determination of reference intervals for 10 serum proteins measured by rate nephelometry, taking into consideration different sample groups and different distribution functions. *J Clin Chem Clin Biochem* 1985;23:157-66.
560. Adam A, Albert A, Calay G, Closset J, Damas J, Franchimont P. Human kininogens of low and high molecular mass: quantification by radioimmunoassay and determination of reference values. *Clin Chem* 1985;31:423-6.
561. Rodger RS, Laker MF, Fletcher K, White TF, Heaton A, Ward MK, Kerr DN. Factors influencing normal reference intervals for creatinine, urea, and electrolytes in plasma, as measured with a Beckman Astra 8 Analyzer. *Clin Chem* 1985;31:292-5.
562. Lamers KJ, Doesburg WH, Gabreels FJ, Romsom AC, Renier WO, Wevers RA, Lemmens WA. Reference values of blood components related to fuel metabolism in children after an overnight fast. *Clin Chim Acta* 1985;145:17-26.
563. van Assendelft OW. Reference values for the total and differential leukocyte count. *Blood Cells* 1985;11:77-96.
564. Cugini P, Murano G, Lucia P, Letizia C, Scavo D, Halberg F, Cornelissen G, Sothorn RB. Circadian rhythms of plasma renin activity and aldosterone: changes related to age, sex, recumbency and sodium restriction. Chronobiologic specification for reference values. *Chronobiol Int* 1985;2:267-76.
565. Gerver WJ, vd Laan CG, Drayer NM, Schaafsma W. Smoothing techniques for obtaining reference values for measurements of children. *Int J Biomed Comput* 1985;16:29-34.

566. Mahr G, Knittel B, Lorunser E, Neyer U, Risch GM. [Amylase and lipase as reference values for the differential diagnosis of chronic kidney failure and pancreatitis]. *Acta Med Austriaca* 1985;12:51-7. [Alemany]
567. Ferro V, Savoia MR, Esposito S, De Vita N, Svelto S. [Contribution to the definition of reference values of various enzymes in the cerebrospinal fluid]. *Quad Sclavo Diagn* 1984;20:379-90. [Italià]
568. Aine E, Morsky P. Lysozyme concentration in tears--assessment of reference values in normal subjects. *Acta Ophthalmol* 1984;62:932-8.
569. Liappis N, Schneider A. [Reference values of sodium, potassium, chloride, calcium, inorganic phosphate and magnesium levels in the cerebrospinal fluid of children]. *Klin Padiatr* 1984;196:370-4. [Alemany]
570. Liappis N, Reimnitz P. [Reference values of sodium, potassium, calcium, chloride and inorganic phosphate excretion in 24-hour urine of healthy children]. *Klin Padiatr* 1984;196:367-9. [Alemany]
571. Kruse K, Kracht U, Kruse U. Reference values for urinary calcium excretion and screening for hypercalciuria in children and adolescents. *Eur J Pediatr* 1984;143:25-31.
572. Klingmuller V, Bergmann E, Claassen P. [Urinary alanine aminopeptidase excretion in children: reference values and effect of nephrotropic contrast media]. *J Clin Chem Clin Biochem* 1984;22:787-9. [Alemany]
573. Quanjor PH. Reference values for adolescents from ages 15 to 20. *Schweiz Med Wochenschr* 1984;114:1507. [Sense resum]
574. Minoia C, Biscaldi GP, Terzi R, Colli M, Marcaletti G, Tempini G. [Reference values of chromium urine in the population residing in Lombardy]. *G Ital Med Lav* 1984;6:169-76. [Italià]
575. Costongs GM, Janson PC, Brombacher PJ. Effects of biological and analytical variations on the appropriate use of "reference intervals" in clinical chemistry. Proposal of a scheme for longitudinal assessment of laboratory values. *J Clin Chem Clin Biochem* 1984;22:613-21.
576. Mortensen H, Molsted-Pedersen L, Schmolker L, Olesen H. Reference intervals for urinary glucose in pregnancy. *Scand J Clin Lab Invest* 1984;44:409-12.
577. Cugini P, Centanni M, Lucia P, Murano G, Letizia C, Scavo D, Bologna U, Biagi L, Dino N, Dolciotti G, et al. [Parametric methods applied to the standardization of gestational reference intervals for blood renin activity, aldosterone and cortisol using the transversal and longitudinal approach]. *Quad Sclavo Diagn* 1984;20:245-57. [Italià]
578. Lakhani S, Jansen AA, Lacko E, Sequeira E. Suggested reference values for serum total protein and fractions of African and Indian middle-class urban women and newborns. *East Afr Med J* 1984;61:606-10. [Sense resum]
579. Schrijver J, Van Rijn JM, Schreurs WH. Re-evaluation of the haptoglobin reference values with the radial immunodiffusion technique. *Clin Biochem* 1984;17:258-62.
580. Zanella A. Reference values for plasma lipids and lipoproteins. *JAMA* 1984;252:343-4. [Sense resum]
581. Marcovina S, Murone M, Dati F, Pfeil H. Reference values for alpha-amylase activity in serum and urine by a new chromogenic amylase method. *Ric Clin Lab* 1984;14:439-42. [Sense resum]
582. Cohen NA. Reference values for creatine kinase MB. *Clin Chem* 1984;30:1102. [Sense resum]
583. [No authors listed] International Federation of Clinical Chemistry (IFCC). Scientific Committee, Clinical Section. Expert Panel on Theory of Reference Values (EPTRV). IFCC Document (1982) stage 2, draft 2, 1983-10-07 with a proposal for an IFCC recommendation. The theory of reference values. Part 2. Selection of individuals for the production of reference values. *Clin Chim Acta* 1984 May 30;139:205F-213F. [Sense resum]
584. Gruemer HD, Miller WG, Chinchilli VM, Leshner RT, Hassler CR, Blasco PA, Nance WE, Goldsmith BM. Are reference limits for serum creatine kinase valid in detection of the carrier state for Duchenne muscular dystrophy? *Clin Chem* 1984;30:724-30.

585. Ojwang PJ, Ogada T, Maina FW, Sekadde-Kigundu CB, Mati JK. Reference values for serum lipids and lipoprotein cholesterol in adults and cord blood of Kenyan Africans. *East Afr Med J* 1984;61:367-71. [Sense resum]
586. Brechbuhler T, Kaeslin M, Wyler F. Reference values of various blood constituents in young minipigs. *J Clin Chem Clin Biochem* 1984;22:301-4.
587. Galvan Guerra E, Ponce de Leon S, Loria A, Robles-Diaz G. [Reference values for amylase in a sample of the Mexican population. Variations according to age]. *Rev Invest Clin* 1984;36:109-14. [Espanyol] [Sense resum]
588. Statland BE, Winkel P. Reference values: are they useful? *Clin Lab Med* 1984;4:61-70.
589. Gomez P, Coca C, Vargas C, Acebillo J, Martínez A. Normal reference-intervals for 20 biochemical variables in healthy infants, children, and adolescents. *Clin Chem* 1984;30:407-12.
590. Alvarez C, Orejas A, Gonzalez S, Diaz R, Colomo LF. Reference intervals for serum lipids, lipoproteins, and apoproteins in the elderly. *Clin Chem* 1984;30:404-6.
591. Valenzuela C. [Sociogenetic reference limits for public health studies in Chile]. *Rev Chil Pediatr* 1984;55:123-7. [Espanyol] [Sense resum]
592. PetitClerc C, Wilding P. International Federation of Clinical Chemistry (IFCC), Scientific Committee, Clinical Section. The theory of reference values. Part 2. Selection of individuals for the production of reference values. *J Clin Chem Clin Biochem* 1984;22:203-8. [Sense resum]
593. Forsberg P, Henriksson A, Link H, Ohman S. Reference values for CSF-IgM, CSF-IgM/S-IgM ratio and IgM index, and its application to patients with multiple sclerosis and aseptic meningoencephalitis. *Scand J Clin Lab Invest* 1984;44:7-12.
594. Lambert D, Daubrosse E, Steinmetz J, Siest G, Debry G. [Reference values of plasma and lipoprotein cholesterol and triacylglycerols in healthy children]. *Arch Fr Pediatr* 1984;41:145-50. [Francès]
595. Lakhani S, Jansen AA, Lacko E, Sequeira E. Suggested reference values for serum total protein and fractions of African and Indian middle-class urban women and newborns. *East Afr Med J* 1984;61:129-32. [Sense resum]
596. Erfurth EM, Norden NE, Hedner P, Nilsson A, Ek L. Normal reference interval for thyrotropin response to thyroliberin: dependence on age, sex, free thyroxin index, and basal concentrations of thyrotropin. *Clin Chem* 1984;30:196-9.
597. West DW, Ash O. Adult reference intervals for 12 chemistry analytes: influences of age and sex. *Am J Clin Pathol* 1984;81:71-6.
598. Stoddard L, Dennis W, Parvin RM, van Assendelft OW. Freeze/thaw stability of transferrin, and reference values obtained with kinetic nephelometry. *Clin Chem* 1984;30:114-5.
599. Toure M, Seck I, Cisse F. [Reference values of glycosylated hemoglobins in black Senegalese]. *Dakar Med* 1984;29:329-34. [Francès] [Sense resum]
600. Toure M, Seck I, Cisse F. [Reference values of HDL cholesterol in healthy Senegalese]. *Dakar Med* 1984;29:117-25. [Francès] [Sense resum]
601. Willroth PO, Zimmermann G. [Reference values for the inhomogeneity parameter V2/V1]. *Z Erkr Atmungsorgane* 1984;163:241-4. [Alemany]
602. Chio LF, Vellayappan K, Tan IK, Chua KS, Wong HB. Reference values for 24 biochemical constituents of children in Singapore. *Ann Acad Med Singapore* 1984;13:85-90.
603. Berg B, Petersohn L, Helm G, Tryding N. Reference values for serum components in pregnant women. *Acta Obstet Gynecol Scand* 1984;63:583-6.
604. Labeeuw M, Pozet N, Gerbault C, Zech P, Traeger J. [Diagnosis of relative hypomagnesurea in calcium lithiasis: reference values for a new parameter]. *Nephrologie* 1984;5:101-5. [Francès]

605. Franck C, Patricot MC, Mathian B, Revol A. [Assay of free urinary cortisol by inverted phase liquid chromatography and radiocompetition. Reference values in adults]. *Ann Biol Clin* 1984;42:221-5. [Francès]
606. Forest JC, Garrido-Russo M, Lemay A, Carrier R, Dube JL. Reference values for the oral glucose tolerance test at each trimester of pregnancy. *Am J Clin Pathol* 1983;80:828-31.
607. Kunze D. Reference values and tracking of blood lipid levels in childhood. *Prev Med* 1983;12:806-9. [Sense resum]
608. Solberg HE. The theory of reference values Part 5. Statistical treatment of collected reference values. Determination of reference limits. *J Clin Chem Clin Biochem* 1983;21:749-60. [Sense resum]
609. Rifkind BM, Segal P. Lipid Research Clinics Program reference values for hyperlipidemia and hypolipidemia. *JAMA* 1983;250:1869-72.
610. Viriyayudhakorn S, Mitarnun W, Pornpatkul M, Jenjareontham R, Khoprasert B, Ruangrairatanaroje P, Musigavon P, Tiewpiyakul N, Keowharmkar W. Reference intervals of twenty-one biochemical constituents of the population in southern Thailand. *J Med Assoc Thai* 1983;66:579-89. [Sense resum]
611. Gaidulis L, Muensch HA, Maslow WC, Borer WZ. Optimizing reference values for the measurement of alpha 1-antitrypsin in serum: comparison of three methods. *Clin Chem* 1983;29:1838-40.
612. Hitz J, Steinmetz J, Siest G. Plasma lecithin:cholesterol acyltransferase -- reference values and effects of xenobiotics. *Clin Chim Acta* 1983;133:85-96.
613. Giacalone G, Morganti G, Monaci E, Bini D, Matini C. [Reference values for the serum concentration of apolipoproteins A and B, the A/B ratio and cholesterol-HDL. Evaluation in a clinically healthy population in Amiata (Grosseto)]. *Quad Sclavo Diagn* 1983;19:377-83. [Italià]
614. Lindstedt G, Eden S, Jagenburg R, Lundberg PA, Mellstrom D, Oden A, Svanborg A. Factors influencing serum free T4 in 70-year-old men. Implications for reference intervals in the elderly. *Scand J Clin Lab Invest* 1983;43:401-13.
615. Seiler D, Nagel D, Tritschler W, Looser S. [Acid phosphatase in blood (substrate: alpha-naphthylphosphate): reference values and diagnostic significance]. *J Clin Chem Clin Biochem* 1983;21:519-25. [Alemany]
616. Witt I, Tritschler W. [alpha 2-Macroglobulin in serum and plasma: reference values with carbobenzoxy-valyl-glycyl-arginine-p-nitroanilide, a chromogenic substrate]. *J Clin Chem Clin Biochem* 1983;21:429-36. [Alemany]
617. Gillard BK, Simbala JA, Goodglick L. Reference intervals for amylase isoenzymes in serum and plasma of infants and children. *Clin Chem* 1983;29:1119-23.
618. Nexø E. Variation with age of reference values for P-cobalamins. *Scand J Haematol* 1983;30:430-2.
619. Ash KO, Clark SJ, Sandberg LB, Hunter E, Woodward SC. The influences of sample distribution and age on reference intervals for adult males. *Am J Clin Pathol* 1983;79:574-81.
620. Grisler R, Ferrari L, Ronco G, Rusconi R, Biella C, Carnelli V. [Reference values of urinary delta-aminolevulinic acid in children]. *Pediatr Med Chir* 1983;5:87-9. [Italià]
621. Nelson W, Cornelissen G, Hinkley D, Bingham C, Halberg F. Construction of rhythm-specified reference intervals and regions, with emphasis on 'hybrid' data, illustrated for plasma cortisol. *Chronobiologia* 1983;10:179-93.
622. Schiele F, Henny J, Hitz J, Petittlerc C, Gueguen R, Siest G. Total bone and liver alkaline phosphatases in plasma: biological variations and reference limits. *Clin Chem* 1983;29:634-41.
623. Revington M. Haematology of the racing Thoroughbred in Australia 1: reference values and the effect of excitement. *Equine Vet J* 1983;15:141-4.
624. Bezemer PD, Netelenbos JC, Mulder C, Theune JA, Stamhuis IH, Straub JP. Determining reference ('normal') limits in medicine: an application. *Stat Med* 1983;2:191-8.

625. Cross JP, Heyns AD. Haematological reference values for the Basotho. *S Afr Med J* 1983;63:480-3.
626. Knobber D, Romahn A, Liappis N. [Concentration levels of thyroxine binding globulin and thyroxine in the blood of euthyroid children. Reliability and diagnostic value of the T4/TBG quotiens--establishing reference values]. *Klin Padiatr* 1983;195:103-6. [Alemany]
627. Liappis N, Rossmoller M, Dorner K, Hildenbrand G. [Reference values of IgA, IgG, IgM, alpha2-macroglobulin, alpha-1- antitrypsin, complement C3, complement C4, ceruloplasmin, haptoglobin, properdin factor B, acid alpha-1 glycoprotein and transferrin concentration in the blood of healthy children]. *Klin Padiatr* 1983;195:107-16. [Alemany] [Sense resum]
628. Albert A, Gueguen R, Sachs C. Treatment of collected reference values and reference interval determination (document H, stage 3, version 1). *Ann Biol Clin* 1983;41:63-79. [Sense resum]
629. Demaurex A, Kung HL, Wirz H. [Creatinine and urine density as reference values in biological monitoring]. *Soz Praventivmed* 1983;28:88-94. [Alemany]
630. Ito R, Statland BE, Sher G, Knutzen V. Assaying plasma prekallikrein (PPK) by a chromogenic based method: analytical considerations and reference values in healthy adults, the pregnant woman, and the neonate. *Adv Exp Med Biol* 1983;156:205-23.
631. Herbeth B, Henny J, Siest G. [Biological variations and reference values of transferrin, immunoglobulin A and orosomucoid]. *Ann Biol Clin* 1983;41:23-32. [Francès]
632. Urdal P, Urdal K, Stromme JH. Age-related reference intervals for creatine kinase BB in serum. *Scand J Clin Lab Invest* 1982;42:621-5.
633. De Leenheer AP, Lambert WE, Claeys I. All-trans-retinoic acid: measurement of reference values in human serum by high performance liquid chromatography. *J Lipid Res* 1982;23:1362-7.
634. Gupta RC, Jain MR, Chandel HL, Punjabi M, Singh PP. Reference interval for phosphohexose isomerase activity in aqueous humor. *Clin Chem* 1982;28:2178. [Sense resum]
635. Lau HK, Howard SF, Tomczak G, Tan TK. A lower reference interval for hemoglobin A1. *Clin Chem* 1982;28:1822-3. [Sense resum]
636. Witt I, Tritschler W. [alpha 1-antitrypsin: reference values in serum and plasma with benzoyl-D,L-arginine-p-nitroanilide, a chromogenic substrate]. *J Clin Chem Clin Biochem* 1982;20:587-91. [Alemany]
637. Knopp RH, Bergelin RO, Wahl PW, Walden CE, Chapman M, Irvine S. Population-based lipoprotein lipid reference values for pregnant women compared to nonpregnant women classified by sex hormone usage. *Am J Obstet Gynecol* 1982;143:626-37. [Sense resum]
638. Butts WC, Lilje DJ. Clinically significant reference intervals. *Am J Med Technol* 1982;48:587-94. [Sense resum]
639. Albers JJ, Bergelin RO, Adolphson JL, Wahl PW. Population-based reference values for lecithin-cholesterol acyltransferase (LCAT). *Atherosclerosis* 1982;43:369-79.
640. Andreasen TL, Hansen E, Olesen H. Reference intervals for haemoglobins estimated by isoelectric focusing. *Scand J Clin Lab Invest* 1982;42:119-22. [Sense resum]
641. Witt I, Trendelenburg C. [Joint study to establish reference values for clinical chemical parameters in childhood]. *J Clin Chem Clin Biochem* 1982;20:235-42. [Alemany]
642. Dorner K, Gaethke AS. Calculation factors and reference values for quantitative lipoprotein electrophoresis in childhood. *Clin Chim Acta* 1982;119:99-105.
643. Raun NE, Moller BB, Back U, Gad I. On individual reference intervals based on a longitudinal study of plasma proteins and lipids in healthy subjects, and their possible clinical application. *Clin Chem* 1982;28:294-300.
644. Aellig A, Albert A, Blin G, Buret J, Daubrosse E, Drosdowsky M, Favre J, Gouget B, Guize L, Henny J, Petitclerc C, Queralto-Compano JM, Ramon-Bauza F, Siest G, Vernet M. *Societe Francaise de*

- Biologie Clinique. Section of physiopathology. Commission "reference values". Utilisation of reference values. (Document J, stage 3, version 1). *Ann Biol Clin* 1982;40:697-708. [Sense resum]
645. Harm K, Wegener M, Dieckvoss E, Nagel HH, Voigt KD. [Plausibility control of clinical chemistry reference values]. *Med Lab* 1982;35:145-52. [Alemany] [Sense resum]
646. [No authors listed] Standardization of blood specimen collection procedure for reference values. International Committee for Standardization in Haematology (ICSH). *Clin Lab Haematol* 1982;4:83-6.
647. Scott TG. A pilot study of the reference values for the commoner haematological and biochemical parameters in Saudi nationals. *J Clin Pathol* 1982;35:69-73.
648. Jolliff CR, Cost KM, Stivrins PC, Grossman PP, Nolte CR, Franco SM, Fijan KJ, Fletcher LL, Shriner HC. Reference intervals for serum IgG, IgA, IgM, C3, and C4 as determined by rate nephelometry. *Clin Chem* 1982;28:126-8.
649. Feldmann D, Read MH, Rosenau A, Lavolloy B, Aymard P. [Assay of total serum bile acids. Reference values in children]. *Ann Biol Clin* 1982;40:181-5. [Francès]
650. Makino T, Takahara K. Reference values: accurate determination of plasma iron in infants by discrete nebulization in atomic absorption. *Clin Chem* 1981;27:2073. [Sense resum]
651. Allner R, Gaschler E, Herrmann B. [Reference values for lipase activity in the serum. Diagnostic significance of alpha-amylase and lipase in the serum]. *Med Welt* 1981;32:1533-7. [Alemany] [Sense resum]
652. Massmann W. Reference values of renal excretion of fluoride. *J Clin Chem Clin Biochem* 1981;19:1039-41.
653. Albert A. Atypicality indices as reference values for laboratory data. *Am J Clin Pathol* 1981;76:421-5.
654. Granerus G, Aurell M. Reference values for ⁵¹Cr-EDTA clearance as a measure of glomerular filtration rate. *Scand J Clin Lab Invest* 1981;41:611-6.
655. Letellier G. [Reference values for 2 well-defined populations: newborn infants and post-partum women]. *Clin Biochem* 1981;14:169-72. [Francès]
656. Weisshaar HD, Sudhoff H, Koller PU, Willamowski KD. [Reference values for a new alpha-amylase determination in serum and urine]. *Dtsch Med Wochenschr* 1981;106:936-9. [Alemany]
657. Sapigni T, Astolfi G, Cavallini L, Cremonini F. [Influence of age on blood glucose levels: percentile reference intervals determined on ambulatory patients]. *Boll Soc Ital Biol Sper* 1981;57:1254-60. [Italià]
658. White GL Jr, Nelson JA, Pedersen DM, Ash KO. Fasting and gender (and altitude?) influence reference intervals for serum bilirubin in healthy adults. *Clin Chem* 1981;27:1140-2. [Sense resum]
659. Karlsen RL, Sterri S, Lyngaas S, Fonnum F. Reference values for erythrocyte acetylcholinesterase and plasma cholinesterase activities in children, implications for organophosphate intoxication. *Scand J Clin Lab Invest* 1981;41:301-2.
660. Hoes MJ, Kreutzer EK, Sijben N. Xanthurenic acid excretion in urine after oral intake of 5 grams L-tryptophan by healthy volunteers: standardisation of the reference values. *J Clin Chem Clin Biochem* 1981;19:259-64.
661. Bienvenu J, Sann L, Bienvenu F, Lahet C, Divry P, Cotte J, Bethenod M. Laser nephelometry of orosomucoid in serum of newborns: reference intervals and relation to bacterial infections. *Clin Chem* 1981;27:721-6.
662. Posma FD, Groenewold H, Kluff O, Tuynman FH. Reference values and analytical performance of the hydroxyproline/creatinine ratio in early morning urine samples. *J Clin Chem Clin Biochem* 1981;19:209-12.
663. Hesse R, Tritschler W, Castelfranchi G, Bablok W. [Antithrombin III: reference values found with a chromogenic substrate (chromozym TH)]. *Blut* 1981;42:227-34. [Alemany]

664. Weisshaar HD, Sudhoff H, Koller PU, Bablok W. [Pancreatic lipase: reference values at 25 degrees C]. *Dtsch Med Wochenschr* 1981;106:239-41. [Alemany]
665. Speich M, Bousquet B, Nicolas G. Reference values for ionized, complexed, and protein-bound plasma magnesium in men and women. *Clin Chem* 1981;27:246-8.
666. [No authors listed] The theory of reference values: international committee for standardization in haematology (ICSH). *Clin Lab Haematol* 1981;3:369-73. [Sense resum]
667. Siest G, Henny J. Commission on "Reference Values". Production of reference values in healthy subjects (document G, stage 3, version 1). *Ann Biol Clin* 1981;39:235-44. [Sense resum]
668. Alvarez X, Piedras J, Cordova MS, López X, Cano R. [Reference values for serum ferritin in Mexican adults, men and women]. *Rev Invest Clin* 1981;33:13-6. [Espanyol] [Sense resum]
669. Bladier D, Perret G, Pre J, Cornillot P. [Interest of the determination of reference values for the monosaccharide composition of the human erythrocyte membrane]. *Ann Biol Clin* 1981;39:109-13. [Francès]
670. Gergely T, Endler AT, Bauer K, Junker E, Schnack H, Abel-Telkes B, Burger E. [HDL-cholesterol reference values in an urban population (Vienna)]. *Med Lab* 1981;34:15-9. [Alemany] [Sense resum]
671. Ulmer HV. [Work-load and work-time with reference to limits of tolerance in physical work]. *Zentralbl Arbeitsmed Arbeitsschutz Prophyl Ergonomie* 1981;31:14-6. [Alemany] [Sense resum]
672. Dybkaer R, Lauritzen M, Krakauer R. Relative reference values for clinical chemical and haematological quantities in 'healthy' elderly people. *Acta Med Scand* 1981;209:1-9.
673. Harris EK. Statistical aspects of reference values in clinical pathology. *Prog Clin Pathol* 1981;8:45-66. [Revisió] [Sense resum]
674. Boyadjian H. Reference values of antithrombin III and plasminogen and their distribution patterns. *Folia Med* 1981;23:5-10. [Sense resum]
675. Giorno R, Clifford JH, Beverly S, Rossing RG. Hematology reference values. Analysis by different statistical technics and variations with age and sex. *Am J Clin Pathol* 1980;74:765-70.
676. Bright M, Wagman E, Shastri S, Nevins M. Race-related differences in reference intervals for creatine kinase. *Clin Chem* 1980;26:1928-8. [Sense resum]
677. Bimenya GS, Lutalo-Bosa AJ, Nzaro E. Serum zinc reference values in healthy Ugandans (HbAA). *East Afr Med J* 1980;57:822-4. [Sense resum]
678. Naus AJ, Borst A, Kuppens PS. The use of patient data for the calculation of reference values for some haematological parameters. *J Clin Chem Clin Biochem* 1980;18:621-5.
679. Seamonds B, Anderson K, Whitaker B. Reference intervals for ferritin: age dependence. *Clin Chem* 1980;26:1515-6. [Sense resum]
680. Kach O, Sturzenegger MH, Zbinden MA, Bucher U, Beck EA. [Normal reference values in frequently carried out laboratory analyses hematologic cytology and blood coagulation)]. *Schweiz Med Wochenschr* 1980;110:1244-9. [Alemany]
681. Kupke IR, Tritschler W, Kather B, Bablok W. [Creatine kinase "NAC-activated": reference values for children]. *Klin Padiatr* 1980;192:348-50. [Alemany]
682. Jernigan JA, Gudat JC, Blake JL, Bowen L, Lezotte DC. Reference values for blood findings in relatively fit elderly persons. *J Am Geriatr Soc* 1980;28:308-14.
683. Scholler R, Avigdor R, Scholler D. [Reference values of steroids and protein hormones during pregnancy]. *Pathol Biol* 1980;28:359-60. [Francès] [Sense resum]
684. van der Sluijs Veer G. [Reference values for lactate dehydrogenase in the serum during childhood and puberty]. *J Clin Chem Clin Biochem* 1980;18:305-6. [Alemany]

685. Boeckx RL, Iosefsohn M, Hicks JM. Reference values for cerebrospinal fluid glutamine concentration in infants. *Clin Chem* 1980;26:601-3.
686. Bavazzano P, Benassi S, Forzieri R, Petrioli G. [Identification of the reference values of serum levels of aspartate and alanine aminotransferases found in 1262 apprentices undergoing examinations for fitness for work]. *Quad Scavo Diagn* 1980;16:100-18. [Italià] [Sense resum]
687. Plauchu H, Claustrat B, Betend B, David M, Francois R. [Synthetic LHRH hypothalamic hormone test in healthy children from birth to adulthood. Reference values]. *Pediatric* 1980;35:110-31. [Francès] [Sense resum]
688. Felding P, Tryding N, Hyltoft Petersen P, Horder M. Effects of posture on concentrations of blood constituents in healthy adults: practical application of blood specimen collection procedures recommended by the Scandinavian Committee on Reference Values. *Scand J Clin Lab Invest* 1980;40:615-21.
689. Plebani M, Bonvicini P, De Besi T, Giordano A, Pesarin F, Ceriotti G. Reference values for alanine and aspartate aminotransferase (ALT and AST) optimized by addition of pyridoxal phosphate. *Enzyme* 1980;25:346-52.
690. Drosdowsky M, Sachs C. Factors to be taken into consideration for blood sampling in view of establishing reference values (document E. Stage 3). *Ann Biol Clin* 1980;38:251-65. [Sense resum]
691. Larsen LH, Wilson M. Age and sex specific reference limits for carbohydrate tolerance. *Med J Aust* 1979;2:674-7.
692. Kristensen SR, Horder M, Pedersen GT. Reference values for six enzymes in plasma from newborns and women at delivery. *Scand J Clin Lab Invest* 1979;39:777-84.
693. Borner U, Szasz G, Bablok W, Busch EW. [A specific fully enzymatic method for creatinine: reference values in serum]. *J Clin Chem Clin Biochem* 1979;17:679-82. [Alemany]
694. Alvarez L, de Miguel E, Calvet E, Ramon F. [Reference values of A.S.T., L.D.H., isoenzymes of L.D.H. and A.D.A. in cerebrospinal fluid of children]. *An Esp Pediatr* 1979;12:779-84. [Espanyol]
695. Colon EJ, de Weerd JP, Notermans SL, de Graaf R. EEG spectra in children aged 8, 9 and 10 years. Reference values. *J Neurol* 1979;221:263-8.
696. Chio LF, Vellayappan K, Tan IK, Wong HB. Reference values for serum immunoglobulins and complement C3 and C4 of Singapore children. *J Singapore Paediatr Soc* 1979;21:210-7. [Sense resum]
697. Rajamaki A, Irjala K, Aitio A. Immunochemical determination of serum transferrin. Reference values, correlation with serum total iron-binding capacity and value in the diagnosis of iron deficiency anaemia and anaemia of chronic disorders. *Scand J Haematol* 1979;23:227-31.
698. Dorner K, Stingl EM, Toeller W, Simon C. [Reference values of lipid metabolism in childhood]. *Monatsschr Kinderheilkd* 1979;127:511-4. [Alemany]
699. Grasbeck R, Siest G, Wilding P, Williams GZ, Whitehead TP. Provisional recommendation on the theory of reference values (1978). Part 1. The concept of reference values. *Clin Chem* 1979;25:1506-8. [Sense resum]
700. Manroe BL, Weinberg AG, Rosenfeld CR, Browne R. The neonatal blood count in health and disease. I. Reference values for neutrophilic cells. *J Pediatr* 1979;95:89-98.
701. Rossing RG, Hatcher WE 3rd. Percentiles as reference values for laboratory data. *Am J Clin Pathol* 1979;72:94-7.
702. Steinmetz J, Panek E, Siest G, Gueguen R. Factors affecting the concentration of triacylglycerols (triglycerides in plasma: reference values for adults. *Clin Chem* 1979;25:924-32.
703. Salvati AM, Camagna A, Samoggia P, Tentori L. Reference values in haematology. A survey in Italy: report from Latium. *Haematologica* 1979;64:296-308. [Sense resum]

704. Parvy P, Huang Y, Kamoun P. Age related reference values for urinary free amino acids: a simple method of evaluation. *J Clin Chem Clin Biochem* 1979;17:205-10.
705. Freer DE, Statland BE, Johnson M, Felton H. Reference values for selected enzyme activities and protein concentrations in serum and plasma derived from cord-blood specimens. *Clin Chem* 1979;25:565-9. [Sense resum]
706. Feldt-Rasmussen U, Hyltoft Petersen P, Date J. Sex and age correlated reference values of serum thyroglobulin measured by a modified radioimmunoassay. *Acta Endocrinol* 1979;90:440-50.
707. Chemnitz G, Schmidt E, Koller PU, Busch EW. [Creatine kinase: reference values and clinical aspects of the revised standard method]. *Dtsch Med Wochenschr* 1979;104:257-60. [Alemany]
708. Graubaum HJ, Wagenknecht C, Meurer W, Stepan J. [Reference values for alkaline phosphatase isoenzymes, GOT isoenzymes and creatinine - investigation of dietary effects]. *Z Med Lab Diagn* 1979;20:24-31. [Alemany] [Sense resum]
709. Butrimovitz GP, Purdy WC. Resolution of age-dependent reference intervals: polynomial regression methodology with applicability to plasma zinc levels in a childhood population. *Clin Biochem* 1979;12:33-6.
710. Fogstrup J, With TK. Urinary total porphyrins by ion exchange analysis: reference values for the normal range and remarks on preformed porphyrins in acute porphyria urine. *J Clin Pathol* 1979;32:109-12.
711. Venturini M, De Grandi R, Calafati P, Conti G, Repetto M, Corciulo G. [Determination of reference values of serum cholesterol]. *Arch Maragliano Patol Clin* 1979;35:145-52. [Italià] [Sense resum]
712. [No authors listed] [Commission on "Proteins". Reference values of serum IgG, IgA and IgM in children from birth to age 15 years (document A, stage 3)]. *Ann Biol Clin* 1979;37:127-34. [Francès] [Sense resum]
713. Gardner MD, Scott R. Frequency distribution and 'reference values' of plasma alkaline phosphatase (EC 3.1.3.1) activity in the adult population of a Scottish new town. *J Clin Pathol* 1978;31:1202-6.
714. Ahonen A, Myllyla VV, Hokkanen E. Measurement of reference values for certain proteins in cerebrospinal fluid. *Acta Neurol Scand* 1978;58:358-65.
715. Heimsoth VH, Graffe-Achelis C, Banauch D, Vollmar J. [Reference values for the urinary glucose concentration test in the adult]. *Med Lab* 1978;31:236-40. [Alemany] [Sense resum]
716. Breebaart K, Becker H, Jongebloed FA. Investigation of reference values of components of cerebrospinal fluid. *J Clin Chem Clin Biochem* 1978;16:561-5.
717. Harwood SJ, Cole GW. Reference values based on hospital admission laboratory data. *JAMA* 1978;240:270-4.
718. Halberg F, Lee JK, Nelson W. Time-qualified reference intervals - chronodesms. *Experientia* 1978;34:713-6.
719. Picard RA. Effect of posture of subject on reference values. *Clin Chem* 1978;24:1083-4. [Sense resum]
720. Delwaide PA. [Current concepts concerning reference values in clinical chemistry]. *Rev Med Liege* 1978 May 1;33:309-26. [Francès] [Sense resum]
721. Davis JR, Avram MJ. Sex-related differences in the reference values for erythrocytic delta-aminolevulinic acid dehydratase activity. *Clin Chem* 1978;24:726-7. [Sense resum]
722. Antonozzi I, Del Castello PG, Morisi G, Ceccarelli P. [Screening for aminoacidopathies in newborns by means of an aminoacid analyzer. Reference values and statistical determinations]. *Ann Ist Super Sanita* 1978;14:781-91. [Italià]
723. Lehmann I, Blatterlein E, Wagenknecht C, Schirgel B. [Reference values of lactate dehydrogenase (LDH), aspartate aminotransferase (GOT) and Aldolase isoenzymes in the plasma of children]. *Padiatr Grenzgeb* 1978;17:245-50. [Alemany] [Sense resum]

724. Siest G, Petitclerc C. Selection of people for the production of population reference values [proceedings] *Ann Biol Clin* 1978;36:217-20. [Sense resum]
725. Grasbeck R, Dybkaer R, Winkel P. Relating observed values to reference values [proceedings] *Ann Biol Clin* 1978;36:193-4. [Sense resum]
726. Cherian AG, Hill JG. Percentile estimates of reference values for fourteen chemical constituents in sera of children and adolescents. *Am J Clin Pathol* 1978;69:24-31.
727. Tibbling G, Link H, Ohman S. Principles of albumin and IgG analyses in neurological disorders. I. Establishment of reference values. *Scand J Clin Lab Invest* 1977;37:385-90. [Sense resum]
728. Coombes EJ, Shakespeare PG, Batstone GF. Age and sex related reference values for serum ribonuclease. *Clin Chim Acta* 1977;79:271-5. [Sense resum]
729. Buffone GJ, Lewis SA. Effect of analytical factors on immunochemical reference limits for complement component C3 in serum of a reference pediatric population. *Clin Chem* 1977;23:994-9.
730. Schiele F, Guilmin AM, Detienne H, Siest G. Gamma-glutamyltransferase activity in plasma: statistical distributions, individual variations, and reference intervals. *Clin Chem* 1977;23:1023-8.
731. Pohlandt F. [Reference values for plasma amino acid concentration in newborn infants]. *Monatsschr Kinderheilkd* 1977;125:441-3. [Alemany] [Sense resum]
732. Nadeau A. [Reference values for blood sugar and insulin during the prolonged fasting test in a Quebec population]. *Union Med Can* 1977;106:343-6. [Francès] [Sense resum]
733. Try K. Reference values for fibrinogen concentration and inhibitors of fibrinolysis. *Scand J Clin Lab Invest* 1977;37:89-90.
734. Vogel J, Petro W. [Reply to "H. Kleinau: Relationship between reference values of arterial and atmospheric PO₂]? *Z Erkr Atmungsorgane* 1977;148:257-8. [Alemany] [Sense resum]
735. Barth JA. [Relationship between reference values of arterial and atmospheric PO₂]? *Z Erkr Atmungsorgane* 1977;148:255-6. [Alemany] [Sense resum]
736. Grasbeck R. Reference values (formerly called normal values). *Chronobiologia* 1977;4:59-61. [Sense resum]
737. Fiet J, Passa P, Dubos G, Tabuteau F, Dreux C. [Study of the reference values and nyctohemeral cycle of serum iron]. *Ann Biol Clin* 1977;35:305-13. [Francès]
738. Schiele F, Galteau MM, Siest G. [Human plasma lactic dehydrogenase activity: individual variations and reference intervals]. *Ann Biol Clin* 1977;35:379-86. [Francès]
739. Strunge P, Trostmann AF. An epidemiological study of a Danish school population. I. Reference values for serum lipids based on a clinical and biochemical study of 284 girls and 268 boys. *Acta Paediatr Scand* 1976;65:689-93. [Sense resum]
740. Gauvin R, Talbot J, Loïselle JM, Page M, Bergeron L. [Reference values applicable to 12 serum parameters: study carried out on an active population of the Quebec area]. *Union Med Can* 1976;105:1714-7. [Francès] [Sense resum]
741. Harris EK. Some theory of reference values. II. Comparison of some statistical models of intraindividual variation in blood constituents. *Clin Chem* 1976;22:1343-50.
742. Norgaard-Pedersen B, Schultz-Larsen P. Reference interval for maternal serum alpha-fetoprotein. *Clin Genet* 1976;9:374-7.
743. Stepan J, Graubaum HJ, Meurer W, Wagenknecht C. [Isoenzymes of alkaline phosphatase - reference values in young people and effects of protein diet]. *Experientia* 1976;32:832-4. [Alemany]
744. Sunderman FW Jr. Current concepts of "normal values," "reference values," and "discrimination values," in clinical chemistry. *Clin Chem* 1975;21:1873-7.

745. Harris EK. Some theory of reference values. I. Stratified (categorized) normal ranges and a method for following an individual's clinical laboratory values. *Clin Chem* 1975;21:1457-64.
746. Siest G, Schiele F, Galteau MM, Panek E, Steinmetz J, Fagnani F, Gueguen R. Aspartate aminotransferase and alanine aminotransferase activities in plasma: statistical distributions, individual variations, and reference values. *Clin Chem* 1975;21:1077-87.
747. Alstrom T, Grasbeck R, Hjelm M, Skandsen S. Recommendations concerning the collection of reference values in clinical chemistry and activity report by the Committee on Reference Values of the Scandinavian Society for Clinical Chemistry and Clinical Physiology. *Scand J Clin Lab Invest* 1975;144(Suppl.):1-44. [Sense resum]
748. Siest G. [Reference values in biology. Use and special value in preventive medicine]. *Pathol Biol* 1975;23:63-70. [Francès]
749. Galteau MM, Siest G, Boura M. [Does mild exercise affect the reference values of the blood parameters?]. *Clin Chim Acta* 1974;55:353-7. [Francès] [Sense resum]
750. Copeland BE, Shruhan C, Day K, Doherty A. Long-term human reference values based on a specific age range. Report of five years experience. *Z Klin Chem Klin Biochem* 1974;12:252. [Sense resum]
751. Dyerberg J, Hjerne N, Nyman G, Olsen JS. Reference values for cord blood lipid and lipoprotein concentrations. *Acta Paediatr Scand* 1974;63:431-6. [Sense resum]
752. Tschanz C, Dauwalder H, Colombo JP. Plasma alkaline phosphatase. Reference values for total activity, L-phenylalanine inhibited fraction, dependency on age, sex and blood grouping. *Clin Biochem* 1974;7:68-80. [Sense resum]
753. Thefeld W, Hoffmeister H, Busch EW, Koller PU, Vollmar J. [Reference values for the determination of GOT, GPT, and alkaline phosphatase in serum with optimal standard methods]. *Dtsch Med Wochenschr* 1974;99:343-4. [Alemany] [Sense resum]
754. Batt AM, Siest G, Loppinet V, Guerin P, Guerguen R, Floc'h AY. [Drugs and reference values in biology. I. Oral contraceptives]. *Ann Biol Clin* 1974;32:245-56. [Francès] [Sense resum]
755. Klebe JG, Winkel P, Lyngbye J. Urinary oestriol excretion in diabetic pregnancy. The problem of reference intervals. *Acta Endocrinol Suppl* 1974;182:52-6. [Sense resum]
756. Muller G. [Reference values of sodium and potassium blood levels]. *Z Med Labortech* 1974;15:184-6. [Alemany] [Sense resum]
757. Winkel P, Jorgensen K. A method for the computation of age-dependent reference intervals. *Scand J Clin Lab Invest* 1973;31:481-3. [Sense resum]
758. Gerhardt W. Proceedings: Specific determination of glucose-6-phosphate dehydrogenase in erythrocytes using the LKB 8600: extraction procedures, stability studies, and reference values calculated per mean erythrocyte. *Quad Sclavo Diagn* 1973;9:464-71. [Sense resum]
759. Sharpe DM, Wilcock AR, Goldberg DM. Automated kinetic spectrophotometric assays of enzyme activities of human cerebrospinal fluid: methods and reference values. *Clin Chem* 1973;19:240-7. [Sense resum]
760. Launay JM, Dreux C. [Studies on platelet serotonin. I. Fluorimetric determination, reference values, 1st results]. *Ann Biol Clin* 1973;31:387-95. [Francès] [Sense resum]
761. Lestin H, Schirgel B, Depersdorff J, Thiele HJ. [Model for multiphasic screening. 3. Hemoglobin determination for the establishment of reference values and anemia incidence in a rural population]. *Dtsch Gesundheitsw* 1972;27:2312-8. [Alemany] [Sense resum]